

## Remarks

### The Amendments

Claim 1 has been amended to delete the recitations “or a functional equivalent thereof” and “solely.” Claim 1 also has been amended to recite that “the heterologous gene expresses an active pyruvate decarboxylase.” This amendment is supported by Example 10 of the specification, which discloses expression of an active pyruvate decarboxylase from a heterologous gene.

Claims 9, 12, and 13 have been rewritten as independent claims.

Claim 14 has been amended to recite bacterial strain “LN-DP1” in place of “LN-P1.” The amendment corrects an obvious typographical error. See Example 6, which describes the “Production of Strain LN-DP1.” (Page 10, line 22 to page 11, line 5.)

New claims 26-28 recite that the bacterium of claims 9, 12, and 13 is a thermophile. These new claims are supported by originally filed claim 3.

New claims 29-31 recite that the bacterium of claims 9, 12, and 13 further comprises an inactivated lactate dehydrogenase gene. New claims 29-31 are supported by originally filed claim 5.

None of these amendments introduces new matter or requires a new search.

### Rejection of claims 1-13 under 35 U.S.C. §112, second paragraph

Claims 1-13 are rejected under 35 U.S.C. §112, second paragraph as being indefinite. Applicants respectfully traverse.

The Office Action asserts that the recitation “but has solely native alcohol dehydrogenase function” in claim 1 is indefinite. As suggested in the Office Action, this recitation has been replaced by the recitation “has native alcohol dehydrogenase function.”

The Office Action also asserts that the recitation “or a functional equivalent thereof” in claim 1 is indefinite. To advance prosecution, this recitation has been deleted.

Withdrawal of these rejections of claims 1-13 is respectfully requested.

Rejection of claims 1-14 under 35 U.S.C. §112, first paragraph

Claims 1-14 are rejected under 35 U.S.C. §112, first paragraph as lacking both written description and enablement. Applicants respectfully traverse.

Written description rejection of claims 1-14

The Office Action asserts that the broad genus of “a heterologous gene encoding pyruvate decarboxylase (pdc) or a functional equivalent thereof” is not adequately described in the specification. Amended claim 1, however, does not recite this broad genus. To advance prosecution, claim 1 has been amended to recite the genus of “a heterologous gene encoding pyruvate decarboxylase.” The specification adequately describes this genus.

Heterologous genes encoding pdc were well known in the art prior to the filing of the application. A heterologous gene is a gene derived from a different species. (See Merriam Webster’s definition of “heterologous”; Appendix A.) The specification teaches that heterologous pdc genes “may be from *Zymomonas sp*, preferably *Z. mobilis* or may be from yeast e.g. the *S. cerevisiae* pdc 5 gene.” (Page 5, lines 1-2.) The nucleotide sequence of at least ten other pyruvate decarboxylase genes derived from species other than a species of gram-positive bacteria (*i.e.*, heterologous genes) were known in the art prior to the January 6, 2000

effective filing date of the application:

- Accession number X87929 (Appendix B) discloses the nucleotide and amino acid sequence for *Kluyveromyces lactis* pyruvate decarboxylase;
- Accession number X81854 (Appendix C) discloses the nucleotide and amino acid sequence for *Nicotiana tabacum* pyruvate decarboxylase;
- Accession number U71122 (Appendix D) discloses the nucleotide and amino acid sequence for *Arabidopsis thaliana* pyruvate decarboxylase-2;
- Accession number U65927 (Appendix E) discloses the nucleotide and amino acid sequence for *Neospora crassa* pyruvate decarboxylase;
- Accession number X92743 (Appendix F) discloses the nucleotide and amino acid sequence for *Oryza sativa* pyruvate decarboxylase;
- Accession number X59546 (Appendix G) discloses the nucleotide sequence and amino acid sequence for *Zea mays* pyruvate decarboxylase;
- Accession number U75311 (Appendix H) discloses the nucleotide and amino acid sequence for *Pichia stipitis* pyruvate decarboxylase;
- Accession number AF098293 (Appendix I) discloses the nucleotide sequence and amino acid sequence for *Aspergillus oryzae* pyruvate decarboxylase;
- Accession number AI728334 (Appendix J) discloses the nucleotide sequence for *Gossypium hirsutum* pyruvate decarboxylase isozyme 1; and
- Accession number Z54096 (Appendix K) discloses the nucleotide sequence of a cosmid from *Schizosaccharomyces pombe* chromosome 1 that includes the nucleotide and amino acid sequence of pyruvate decarboxylase (page 5).

These sequences were retrieved from the National Center for Biotechnology Information (NCBI) database and bear dates that are at least two and one half months before the priority date of the present application. A specification need not disclose what is well known to those skilled in the art and preferably omits that which is well known to those skilled and already available to the public. *In re Buchner*, 929 F.2d 660, 661 (Fed. Cir. 1991); *Hybritech, Inc. v. Monoclonal Antibodies, Inc.*, 802 F.2d 1367, 1384 (Fed. Cir. 1986). Thus, the teachings of the specification,

together with the knowledge in the art of numerous “heterologous genes encoding pyruvate decarboxylase,” adequately demonstrates that applicants were in possession of the invention of claims 1-13 when the present specification was filed.

New claims 26-31 should also not be subject to this rejection. They each ultimately depend from claim 1 and thus also recite gram-positive bacterium transformed with a heterologous gene encoding *pdh*.

Claim 14 recites particular deposited strains of bacteria. A reference in the specification to a deposit constitutes an adequate description of the deposited material sufficient to comply with the written description requirement. *Enzo Biochem v. Gen-Probe Incorporated*, 296 F.3d 1316, 1325 (Fed. Cir. 2002). Thus claim 14 also is adequately described.

Withdrawal of this rejection of claims 1-14 is respectfully requested.

Enablement rejection of claims 1-14

The Office Action asserts that claims 12-14 are not enabled because they recite novel vectors and strains of microorganisms, some of which have been deposited, but have not been indicated as being publicly available. (Paper 12, page 5, lines 10-16.) Applicants respectfully traverse.

Claim 12 recites a Gram-positive bacterium transformed with a plasmid, pFC1. The specification enables how to make and use plasmid pFC1. The specification discloses that plasmid pFC1 “was formed from a fusion of pAB124 (Bingham et al., *Gen. Microbiol.*, 114, 401-408, 1979) and pUC18.” (Page 12, lines 18-20.) The specification also teaches that as a result of fusing pAB124 and pUC18, plasmid pFC1 has a restriction enzyme cleavage map as shown in Figure 8. The plasmid pAB124 was known and used at the time the application was filed. Bingham (*J. Gen. Microbiol.* 119, 109-115, 1980) teaches, at Figure 2, a restriction

enzyme cleavage map of plasmid pAB124 (Page 114; Appendix L.) The sequence the plasmid pUC18 was known at the time the application was filed. GenBank accession number L09136 discloses the sequence of the complementary strand of pUC18. See Appendix M. One of skill in the art would have been able to make and use plasmid pFC1 as shown in Figure 8 using the well known structures of pAB124 and pUC18 and the routinely practiced techniques of molecular biology.

Claims 13 and 14 recite ten strains of bacteria. The recited bacterial strains are LN, LN-T (31), LN-T (32), TN, TN-P1, TN-P3, LN-S (J8), LN-D, LN-D11, and LN-DP1. Four of the recited bacterial strains, LN, TN, LN-S (J8), and LN-D, have been deposited with the National Collections of Industrial, Food and Marine Bacteria (NCIMB). These strains will also be made publicly available upon issuance of a patent. See the declaration of the inventors that accompanies this response. Appendix N. Thus each of these four bacterial strains are enabled.

The specification teaches one of skill in the art how to make and use the remaining six bacterial strains without undue experimentation. The specification teaches how to make and use bacterial strains TN-P1 and TN-P3. The specification discloses that bacterial strain TN-P1 is produced from bacterial strain TN after transformation with plasmid pBST22-zym. (Example 8; page 11, lines 21-24.) The specification discloses that bacterial strain TN-P3 is produced by transforming bacterial strain TN with plasmid pFC-PDC1. (Example 9, page 12, lines 15-18.) Those of skill in the art routinely practice transformation of bacteria. Thus, one of skill in the art could readily produce bacterial strains TN-P1 and TN-P3 using the deposited TN bacterial strain.

The specification also teaches one of skill in the art how to make and use bacterial strains LN-T (E31, E32). The specification teaches that bacterial strains LN-T (E31, E32) are “spontaneous transposon mutants from strain LN. Both strains are lactate deficient.” (Page 9,

lines 6-7.) Thus strain LN-T (E31, E32) bacteria are strain LN bacteria that have spontaneously lost LDH activity. One of skill in the art would have been able to produce and identify LN-T strain bacteria by merely culturing the deposited LN strain bacteria and screening the cultured bacteria for LDH activity. Methods of assaying LDH activity of bacteria were also well known at the time the application was filed. Payton *et al.* (FEMS Microbiology Letters, 26, 333-336, 1985; Appendix O) teaches the identification of *Bacillus stearothermophilus* that lack LDH activity. Screening for *B. stearothermophilus* that lacked LDH activity “was achieved by selection for resistance to a suicide substrate, fluoropyruvate.” (Page 333, lines 8-9 of the Summary; See also page 334, column 1, line 15 to column 2, line 40.)

The specification also teaches one of skill in the art how to make and use bacterial strain LN-D11 without undue experimentation. The specification discloses that LN-D11 bacteria are produced from strain LN-D bacteria “after repeated subculture.” (Page 10, line 13.) LN-D11 bacteria are similar to strain LN-D bacteria except they are sensitive to kanamycin. See Figure 1, which discloses that LN-D strain bacteria are  $spo^-$ ,  $ldh^-$ ,  $km^R$ , (sporulation deficient, LDH recombination mutant, and kanamycin resistant) and LN-D11 bacteria are  $spo^-$  and  $ldh^-$  (sporulation deficient and LDH recombinant mutant). One of skill in the art, using the well known techniques of cell culture would have been able to subculture the deposited LN-D strain bacteria and screened it for kanamycin resistance without undue experimentation.

The specification also teaches one of skill in the art how to make and use bacterial strain LN-DP1. The specification discloses that LN-DP1 bacteria are “produced from strain LN-D11 after transformation with the replicative plasmid pBST22-zym.” (Page 10, lines 24-25.) One of skill in the art could readily have produced bacterial strain LN-D11 as indicated above. One of skill in the art would further have been able to transform the LN-D11 strain bacteria with

plasmid pBST22-zym by routine experimentation. Thus, one of skill in the art would also have been able make and use bacterial strain LN-DP1 without having to resort to undue experimentation.

Withdrawal of this rejection to claims 12-14 is respectfully requested.

Rejection of claims 1, 2, 8, and 11 under 35 U.S.C. §102(b)

Claims 1, 2, 8, and 11 are rejected under 35 U.S.C. §102(b) as being anticipated by Danilevich *et al.* (*Molecular Biology*, 1994, 28(1):158-166; “Danilevich”). Applicants respectfully traverse.

To reject claims as anticipated, each and every element as set forth in the claim must be found either expressly or inherently described in a single prior art reference. *Verdegaal Bros. v. Union Oil Co. of California*, 814 F.2d 628, 631 (Fed. Cir. 1987). See also MPEP § 2131. Danilevich does not teach each and every element of claims 1, 2, 8, and 11.

Claim 1, the independent claim of the rejected set, is directed to a Gram-positive bacterium that has been transformed with a heterologous gene encoding pyruvate decarboxylase but that has native alcohol dehydrogenase function. Danilevich is cited as teaching “an identical Gram-positive bacterium belonging to the *Bacillus sp.* (Paper 12, page 6, lines 9-10.)

Danilevich teaches the construction of a recombinant plasmid that encodes *pdh*. The plasmid was transformed into a Gram-positive bacterium, *B. subtilis*. *B. subtilis*, however, lacks native alcohol dehydrogenase function. “The *B. subtilis* genome lacks both genes (*pdh* and *adh*).” (Page 105, second column, lines 14-15; See also page 109, first column, lines 29-30.) Thus Danilevich does not expressly or inherently teach a Gram-positive bacterium that has been transformed with a heterologous gene encoding pyruvate decarboxylase and that “has native

alcohol dehydrogenase function.” Danilevich does not expressly or inherently teach each and every element of claim 1 or of dependent claims 2, 8, and 11 and does not anticipate these claims. Similarly, Danilevich does not anticipate new claims 26-31, which depend from claim 1.

Withdrawal of this rejection is respectfully requested.

Rejection of claims 1-8, 10, and 11 under 35 U.S.C. §103(a)

Claims 1-8, 10, and 11 are rejected under 35 U.S.C. §103(a) as being unpatentable over Danilevich in view of Hartley *et al.* (*Biotechnol.*, 1983, 145(1):390-396; “Hartley”) and “the common knowledge in the art of molecular biology.” (Paper 12, page 6, line 24.) Applicants respectfully traverse.

Each of claims 1-8, 10, and 11 recites a Gram-positive bacterium that (1) has been transformed with a heterologous gene encoding pyruvate decarboxylase and (2) has native *adh* function. To reject these claims as *prima facie* obvious the Patent Office must meet three criteria:

First, there must be some suggestion or motivation, either in the references themselves or in the knowledge generally available to one of ordinary skill in the art, to modify the reference or to combine reference teachings. Second, there must be a reasonable expectation of success. Finally, the prior art reference (or references when combined) must teach or suggest all the claim limitations.

MPEP § 2143. The Patent Office has failed to meet the second criterion.

Danilevich is cited as teaching a Gram-positive bacterium with native *adh* activity that has been transformed with a plasmid that comprises a heterologous *pdc* gene. (Claim 1; Paper 12, page 6, lines 4-7.) Danilevich also is cited as teaching that the bacterium is a *Bacillus sp.* (claim 2), a heterologous *pdc* gene from *Z. mobilis* (claim 8), and transformation of the



bacterium with a plasmid comprising the heterologous gene (claim 11). (Paper 12, page 6, lines 9-11.) Hartley is cited as teaching the use of a thermophilic microorganism such as *B. stearothermophilus* for production of ethanol. (Claims 2-4; Paper 12, page 7, lines 14-15.) Hartley also is cited as teaching inactivation of the lactate dehydrogenase gene (claim 5) and a heterologous *pdc* gene from *Zymomonas sp.* or *S. cerevisiae* (claim 7). (Paper 12, page 8, lines 15-21.) The Office Action asserts:

With the reference of Danilevich et al. which teaches the introduction of *Z. mobilis* pyruvate decarboxylase into *B. subtilis* and the reference of Hartley et al. which teaches the advantages of using a thermophilic *Bacillus* which has been genetically modified for alcohol production, it would have been obvious to one of ordinary skill in the art to transform *B. stearothermophilus* instead of *B. subtilis* as taught by Danilevich et al. and develop a strain for production of ethanol. With the common knowledge prevailing in the art of molecular biology, it would have been obvious to use methods in which the heterologous gene is introduced through a plasmid or incorporate it into the chromosome of the host bacterium. One of ordinary skill in the art would have been motivated to do so as Hartley et al. teach that use of thermophilic bacteria for production of ethanol has certain economic advantages over other fermentation methods. One of ordinary skill in the art would have a reasonable expectation of success since decarboxylase of *Z. mobilis* and the reference of Hartley et al. suggest the same and in addition provide methods to obtain lactate dehydrogenase non-producers.

Paper 12, page 8, lines 3-16.

As indicated above, Danilevich teaches that *B. subtilis* was transformed with a plasmid comprising the *pdc* gene. *B. subtilis* does not have native *adh* function. Hartley teaches manipulation of the genome of a thermophilic, Gram-positive bacterium that has a native *adh* gene (*B. stearothermophilus*) to inactivate the *lactate dehydrogenase (ldh)* gene. One of ordinary skill in the art, however, would not have combined these teachings to arrive at a Gram-

positive bacterium which has been transformed with a heterologous gene encoding pyruvate decarboxylase, and that has native alcohol dehydrogenase function.

Danilevich teaches that the metabolism of a bacterium having both *pdc* and *adh* gene function may be unpredictably altered: “Introduction of two actively expressed genes, *pdk* and *adh*, into *B. subtilis* may result in ethanol production or alter the cell metabolism in an unpredictable manner.” (Page 109, second column, lines 33-36, emphasis added.) Thus, one of ordinary skill in the art would not have reasonably expected to be able to produce a bacterium comprising both an active, expressed heterologous *pdc* gene and native *adh* gene function.

Hartley does not remedy the deficiencies of Danilevich. Hartley teaches that “one might *try* a recombinant DNA approach to introduce and express the gene for pyruvate decarboxylase (PDC) from yeast or *Zymomonas* [*i.e.*, cells that have *adh* activity], which gives acetaldehyde + CO<sub>2</sub> from pyruvate (ADH then converts the acetaldehyde to ethanol).” (Page 896, line 41 to page 897, line 2, emphasis added.) But Hartley does not teach or suggest that a bacterium with both *adh* function and an active, expressed heterologous *pdc* gene could be successfully produced. In fact, if one of ordinary skill in the art looked to Danilevich for an expression vector encoding a *pdc* gene, she would have been taught that introduction of *pdk* and *adh* genes into a Gram-positive bacterium could alter its metabolism and would not reasonably have expected success.

In fact, in contrast to what would have been expected from the teachings of the cited art, the claimed bacteria have the unexpected properties of a significant increase in ethanol production and superior growth characteristics.

Expression of *pdc* has resulted in a significant increase in ethanol production by the recombinant organism and has unexpectedly improved the organism's growth characteristics. Recombinant

microorganisms, which prior to transformation with the *pdc* gene were highly unstable and difficult to culture, show significant increases in growth and survival rates both aerobically and anaerobically as well as an increase in the rate of ethanol production near to theoretical yields.

Page 4, lines 4-10.

One of ordinary skill in the art would have expected even less success at producing the bacterium of dependent claim 3. Claim 3 is directed to a thermophilic Gram-positive bacterium which is transformed with a heterologous gene encoding pyruvate decarboxylase that is expressed and active and which has native alcohol dehydrogenase function. Hartley teaches that a suitable vector for expression of *pdc* in a thermophilic bacterium was not yet available and, even if it were available a bacterium transfected with such a vector may not gain *pdc* function: “This strategy depends on developing suitable vectors for the thermophile and might fail if the enzyme were insufficiently thermostable.” (Page 897, lines 2-4.)


Thus, according to Hartley, suitability of the vector is critical. Danilevich does not teach that the disclosed *pdc* expression vector is suitable for *pdc* expression in thermophilic bacteria. Danilevich teaches an expression vector that contains the *pdc* gene for use in Gram-positive bacterium. (Page 106, first column, lines 18-19.) But the vector was only tested in the Gram-positive bacterium *B. subtilis*, which is not a thermophile. Thus, the ordinary artisan would not have known whether use of the expression vector taught by Danilevich could satisfy either of the criteria set forth by Hartley: suitability for expression in thermophiles and stability of the expressed enzyme. Hartley’s caveat regarding suitability of vectors, combined with Danilevich’s teaching that a bacterium transfected with both *adh* and *pdc* genes may have altered metabolism and Danilevich’s lack of teaching that the disclosed vector will express a functional *pdc* protein

in thermophilic bacteria, would not have provided the ordinary artisan with a reasonable expectation of success of producing the thermophilic bacterium recited in claim 3.

Withdrawal of this rejection to claims 1-8, 10, and 11 is respectfully requested.

Respectfully submitted,

Date: February 14, 2003

By:   
Michelle Holmes-Son  
Registration No. 47, 660

BANNER & WITCOFF  
1001 G Street, N.W., Eleventh Floor  
Washington, D.C. 20001-4597  
(202) 508-9100

Amendments to the Claims

C1 1. (Currently Amended) A Gram-positive bacterium which has been transformed with a heterologous gene encoding pyruvate decarboxylase, wherein the heterologous gene expresses an active pyruvate decarboxylase, or a functional equivalent thereof, but and wherein the bacterium has solely native alcohol dehydrogenase function.

2. (Original) A Gram-positive bacterium according to claim 1 wherein the bacterium is a Bacillus sp.

3. (Currently Amended) A Gram-positive bacterium according to claim 1 wherein the bacterium is a thermophile.

C2 4. (Currently Amended) A Gram-positive bacterium according to claim 2 wherein the Bacillus is selected from B. stearothermophilus; B. calvodox; B. caldotenax, B. thermoglucosidasius, B. coagulans, B. licheniformis, B. thermodenitrificans, and B. caldolyticus.

5. (Currently Amended) A Gram-positive bacterium according to claim 1 wherein the gene encoding lactate dehydrogenase expression has been inactivated.

6. (Original) A Gram-positive bacterium according to claim 5 in which the lactate dehydrogenase gene has been inactivated by homologous recombination.

7. (Previously Amended) A Gram-positive bacterium according to claim 1 in which the heterologous gene is from Zymomonas sp or from Saccharomyces cerevisiae.

8. (Original) A Gram-positive bacterium according to claim 7 in which the heterologous gene is from Z. mobilis.

C3 9. (Currently Amended) A Gram-positive ~~bacteria according to claim 7~~ bacterium comprising a native adh gene and which has been transformed with a ~~in which the heterologous gene is pdc 5~~ gene from *S. cerevisiae* *S. cerevisiae*.

10. (Previously Amended) A Gram-positive bacterium according to claim 9 wherein the heterologous gene is incorporated into the chromosome of the bacterium.

11. (Previously Amended) A Gram-positive bacterium according to claim 1 in which the bacterium has been transformed with a plasmid comprising the heterologous gene.

C4 12. (Currently Amended) A Gram-positive bacterium ~~according to claim 11,~~ comprising a native adh gene and which has been transformed with a plasmid comprising a heterologous gene encoding pyruvate decarboxylase, wherein the plasmid is pFC1.

C5 13. (Currently Amended) A Gram-positive ~~bacteria according to claim 1~~ bacterium comprising a native adh gene and which has been transformed with a heterologous gene encoding pyruvate decarboxylase wherein the heterologous gene is operatively linked to the lactate dehydrogenase promoter from Bacillus strain LN (NCIMB accession number 41038).

C6 14. (Currently Amended) Strains LN (NCIMB accession number 41038); LN-T (E31, E32); TN NCIMB accession number 41039); TN-P1; TN-P3; LN-S (J8) (NCIMB accession number 41040); LN-D (NCIMB accession number 41041); LN-D11 and ~~LN-P1~~ LN-DP1.

15-25. (Canceled)

26. (New) The gram-positive bacterium of claim 9 wherein the bacterium is a thermophile.


27. (New) The gram-positive bacterium of claim 12 wherein the bacterium is a thermophile.

28. (New) The gram-positive bacterium of claim 13 wherein the bacteria is a thermophile.

29. (New) The gram-positive bacterium of claim 9 further comprising an inactivated lactate dehydrogenase gene.

30. (New) The gram-positive bacterium of claim 12 further comprising inactivated lactate dehydrogenase gene.

31. (New) The gram-positive bacterium of claim 13 further comprising inactivated lactate dehydrogenase gene.



## Merriam-Webster's COLLEGIATE® DICTIONARY



Download  
Merriam-Webster  
Toolbox  
up words  
on the

- ▶ Home ▶ Help
- ▶ Word of the Day
- ▶ Word Games
- ▶ Word for the Wise
- ▶ Books and CDs
- ▶ Company Info
- ▶ Customer Service
- ▶ Network Options
- ▶ Language Zone
- ▶ The Lighter Side
- ▶ Site Map



- Atlas
- Reverse Dictionary
- Rhyming Dictionary
- Collegiate® Dictionary
- Collegiate® Thesaurus
- Unabridged

One entry found for **heterologous**.

Main Entry: **het·er·ol·o·gous**

Pronunciation: -'ră-l&-g&s

Function: *adjective*

Etymology: *heter-* + *-logous* (as in *homologous*)

Date: 1893

: derived from a different species <*heterologous* DNAs>

<*heterologous* transplants>

- **het·er·ol·o·gous·ly** *adverb*



hetero

STU  
LO

Get the **Top 10 Most Popular Sites for "heterologous"**

For **More Information on "heterologous"** go to [Britannica.com](http://Britannica.com)

### Shopping



NEW! Britannica 2003  
Ultimate Reference  
Suite DVD  
Price: USD \$69.95  
Only \$49.95 after mail-  
in rebate!



Webster's 3rd New  
International  
Dictionary on CD-ROM  
Price: USD \$48.96  
You save 30%!

- ▶ **Search the Unabridged Dictionary on-line**  
and enjoy enhanced versions of Merriam-Webster's Collegiate®  
Dictionary and Thesaurus at Merriam-Webster Unabridged.

- ▶ **There's a word for you . . .**  
In fact, there's a new word waiting in your e-mail in box every morning  
when you subscribe to the Word of the Day.

- ▶ **Listen to Word for the Wise radio programs**  
Listen to broadcasts or read transcripts of previous Word for the Wise  
programs on public radio.



### Pronunciation Symbols

\&\ as a and u in abut

\&\ as e in kitten

\&r\ as ur/er in further

\a\ as a in ash

\A\ as a in ace

\ä\ as in mop

\au\ as u in ut

\ch\ as ch in chin

\e\ as e in bet

\E\ as ea in easy

\g\ as g in go

\i\ as i in hit

\I\ as i in ice

\j\ as j in job

\ng\ as ng in sing

\O\ as o in go

\o\ as aw in law

\oi\ as oy in boy

\th\ as th in thin

\[th]\ as th in the

\ü\ as oo in loot

\u\ as oo in foot

\y\ as y in yet

\zh\ as si in vision





```

VNGFFGSFCEYVPILQVVVLEQSHDLERLIGDVSIFHDVDDPSEIDSCVRTLFWGKR
PVYMGRLRSKDATKLVPSSSLNGNIADKMGIKNTFFQTDTIKRVIDKIIAEVYASSRPL
IVVDALIDRYNYNSTIQNFLTETGIPFVTTLMSSKGSIDESLPNFVGTFLGTMSQPIVR
EYMNADCTLILGCMIEFNKNSYCRFNYKSKNQILLWNRVVKIENNIIPDILLHELLP
QLIASLDTTKIVNSRPVTIPNMI PRVEPQPVTFLRQEYLWFKMSTWLKQGDV I I SESG
TSAIGLLQKQKFPDNTLRVLSQAIWNSSGYSIGACLGILAAAYRDMGTLDKHRIILMVG DG
SLQFTFQELSTILTHGFKPYIFVINNQGYTVDRTLNREKTHLNATYFDIQWELLKLP
SLFYSQEYFKRRCMSVGELNSLLSDKEFNKSDQLKIVELILPSMDVPVLLDPRDDSSD
DESSPQHCRPRT"

```

gene 2517..3008  
 /gene="CYC1"  
misc\_feature 2517..2543  
 /gene="CYC1"  
 /note="CRE1 consensus"  
misc\_feature 2649..2660  
 /gene="CYC1"  
 /note="ROX1 consensus"  
misc\_feature 2710..2721  
 /gene="CYC1"  
 /note="HAP 1 consensus"  
misc\_feature 2799..2904  
 /gene="CYC1"  
 /note="HAP2/3/4/5 consensus"  
CDS 3006..>3008  
 /gene="CYC1"  
 /function="respiratory chain"  
 /codon\_start=1  
 /product="cytochrome C"  
 /protein\_id="CAA61156.1"  
 /db\_xref="GI:4379104"  
 /translation="M"

BASE COUNT 963 a 588 c 549 g 908 t

ORIGIN

```

1  aagttcattt caatgtcttt cggaaaacct atagttctat cctgtcccta atagtgttca
61  tattgggcag tggattcttt ttatttatgt taagcattcc ctatgtacct cactaggaaa
121 agggcacttg gacatcataa ttcttccttt tatagcaata cttttctttc tagcttccgt
181 cctacctttt tcataccgga ttccaccttg tcagttcata ataatatcaa tactcaccta
241 ttaagagcca catgacgtgc gtgcgattga gtacttaaaa actactgtta ccttttgtcc
301 atcgtgtaac tttgcatttt ctttgtcaca ttttctgtga caagaactaa cttgacaaaa
361 ttgattttcc ttgaagagaa gtcgaattaa aagttaaca ggaatatact gggacggaca
421 aaaagatgga aaccaagacc ctgatacact caggggctgc taaggagatg agctatactg
481 agaggtataa tgtcgcaccg ctaatacctt tgccctgagta tctttttcat agattatttc
541 agctcaactg ccgtactgtg tttggagttg ccaattattc aactgcgaaa ctatatcaag
601 ccatagcagt tagcgggata catgggattc aaacgataaa tcaattgaac acatcatttg
661 cagtagatgc atacgggaga gctatcggag tgagctgcta cgtcactagc gaactctgcag
721 aattaggcca tgttaatggt ttttttggat ctttttgtga atatgttccc atcttacagg
781 tagtcgtctt ggaacagtct catgatcttg agaggttgat tggagatgtt tcaatctttc
841 atgacgtagt ggatgatcct tctgagatag acagttgtgt acgaacgctt ttttggggga
901 aacgcctgtt ttatatgggc ctccgatcga aggatgccac gaaacttgct cccagtagtt
961 ctctaaacgg aaacatagca gataagatgg gcataaagaa cactttcttt caaacggaca
1021 caataaagag ggttatagac aaaatcattg ccgaagttaa cgcttcatca aggccattga
1081 tcgtgggtga tgcattaatt gatagatata attacaacag cacaatacaa aatttcctaa
1141 cggaaacagg aataccattt gtgacaacct taatgtcaaa aggttctatt gacgaaagct
1201 taccacactt tgttgggact ttcttaggta ccatgtcaca gccattgtt cgagaatata
1261 tgaataatgc agattgtacg ttaattttag gatgtatgat cgaaaacttc aaaaactcgt
1321 actgtagatt caactacaag agtaaaaacc aaatcttact ttggaatgat agagttaaaa
1381 ttgaaaataa tataatacct gatattttac ttcattgaact actcccacag ctaatagcgt
1441 cattagatac taccaaaata gtaaaactct gtccagtaac aataccgaat atgataccca
1501 gagtagaacc gcaaccggtg acatttttgc gacaggagta cttttggtt aagatgtcta
1561 catggctaaa acaaggtgac gttattattt ctgaatctgg tacctcagct attggtctcc

```

```
1621 tacaacaaaa attcccagat aatactagge tagtatccca agcaatttgg aattcatctg
1681 gttattctat cgggtgcatgc cttggaattt tggcagcata tcgcgatatg gggacattgg
1741 ataagcatag gattatatta atggtgggtg atgggtcttt acaatttact ttccaagaac
1801 taagcaccat attgacgcac gggttcaaac cttacatttt tgaataaat aatcaagggt
1861 ataccgttga cagaaccttg aatagagaaa agacctatct gaacgtaca tacttcgata
1921 tccaaccgtg ggaactatta aaactccgt cacttttcta tccaagag tatttcaaga
1981 gaagatgtat gtcggttggg gaattgaata gtttattaag cgataaagag ttcaataaat
2041 ccgatcagtt gaaaattgta gaactaatcc ttccatccat ggatgttcca gtactactcg
2101 atccacggga tgacagtagt gatgatgaat cttctccgca gcataaaaga ccagaaacgt
2161 gaaaggagaa ggggtgaaaca gatatcagag aactttcacg cttcaaatat atatatatt
2221 taataagcac tgaacacaca agtacctgct tgggtgatcag tctcaatggt aaccgcatta
2281 ccattaacaa cacggcacgc ttttcgtatc ggcactaacg aagagcacac caaaagtcac
2341 gtgtgcatta taccgatatt cagcgaaga acagaatgaa gtaaataaat actctattga
2401 tgtcaaatat agataatatg agcgaaaaaa tatcatcttc tcagggtaac actgatcaga
2461 gatattccag aaccattacc attaatgtaa ttgtcacgtg ccggtcagta cgtatcacgt
2521 gactgtaaca tacatcagcc aaccaatcag atcgcttccg ttcacagtga cgtccaacct
2581 ctgacaagaa ttgcttttcc ccacgcttta atccccatta agattaccct gcagcaaac
2641 gcacaacggg aaaccaatga gaaaaaagtc tcagaggctt cacagaattt ggctcatctc
2701 gtgtcttgac cgatgttccc caaatgatag acttgtttcg aagcattttt tcggtttaag
2761 tagttaaaaa caggtaggag aaaaatttta gtatgcgcac tatggtttgt tgataatttc
2821 atatataact aacatattaa gatttgatat tcgattgaat ttgatgttcc tttagcaagt
2881 ctttttagcg ttttcttttt tctcctttct atcttaatta aagtttccca aactatcagt
2941 attaaggatt gattcgtcaa atcaaataac tgttacacaa ccaacacaac taatataaag
3001 tcataatg
```

//

Revised: July 5, 2002.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Oct 3 2002 17:48:23



PubMed

Nucleotide

Protein

Genome


Structure

PopSet

Taxonomy

CGCTCAGGATACGACTTCGCTCCCTAGAGGATCGGATCCCGGCGATTATATATACCTCGA  
 TTCTCTATATTCGCGGATGGGATATATACACACACGCGCGCGCATAGCATGAC  
 CGCCATGATTCGCGGATGGGATATATACACACACGCGCGCGCATAGCATGAC  
 CACAGACATACGCTTCGCTCCCTAGAGGATCGGATCCCGGCGATTATATATACCTCGA

Nucleotide

Search  for

Limits

Preview/Index

History

Clipboard

Details



default


☐ 1: X81854. *N.tabacum* mRNA fo...[gi:551260]

Links

LOCUS NTMRNAPD1 1255 bp mRNA linear PLN 06-NOV-1995  
 DEFINITION *N.tabacum* mRNA for pyruvate decarboxylase 1.  
 ACCESSION X81854  
 VERSION X81854.1 GI:551260  
 KEYWORDS pyruvate decarboxylase; tobPDC1.  
 SOURCE common tobacco.

ORGANISM *Nicotiana tabacum*

Eukaryota; Viridiplantae; Streptophyta; Embryophyta; Tracheophyta;  
 Spermatophyta; Magnoliophyta; eudicotyledons; core eudicots;  
 Asteridae; euasterids I; Solanales; Solanaceae; *Nicotiana*.

REFERENCE 1 (bases 1 to 1255)

AUTHORS Bucher,M., Brander,K.A., Sbicego,S., Mandel,T. and Kuhlemeier,C.

TITLE Aerobic fermentation in tobacco pollen

JOURNAL Plant Mol. Biol. 28 (4), 739-750 (1995)

MEDLINE 95375236

REFERENCE 2 (bases 1 to 1255)

AUTHORS Mandel,T.

TITLE Direct Submission

JOURNAL Submitted (23-SEP-1994) T. Mandel, Institute of Plant Physiology,  
 Altenbergrain 21, 3013 Bern, FRG

FEATURES Location/Qualifiers

source

1..1255

/organism="Nicotiana tabacum"

/sub\_species="samsun"

/db\_xref="taxon:4097"

/tissue\_type="leaves after anoxic treatment"

/dev\_stage="adult"

gene

1..1255

/gene="tobPDC1"

CDS

&lt;1..&gt;1255

/gene="tobPDC1"

/EC\_number="4.1.1.1"

/codon\_start=3

/product="pyruvate decarboxylase"

/protein\_id="CAA57447.1"

/db\_xref="GI:551261"

/db\_xref="SWISS-PROT:P51845"

/translation="AADGYARARGVGACVVTFTVGGLSVLNAIAGAYSENLP LICIVG

GPNSNDYGTNRILHHTIGLQDFSQEP RCFQTVTCYRAVNNLEDAHELIDTAVSTALK

ESKPVIYISIGCNLPGIPHPTFSREPVPFALSPRLSNMMGLEAAVEAAAEFLNKAVKPV

LVGGPKMRVAKASDAFVELSDACGYAVAVMPSAKGLFPEHHSHFIGTYWGA VSTAFCA

EIVESADAYL FAGPIFNDYSSVGYSLLLKKEKAIIVQPDRVTIGNGPAFGCVLMRDFL

AALAKRLKHNPTAFENYHRIYVPEGHPLKCEPKEALRVNVLFQHIQNMLSGDSV VIAE

TGDSWFNCQKLKLPKGC GYEFQM QYGSIGWSVGATLGYAQA APEKRVIACIGDGSFQV

TAQDISTMLRCGQRTIIFLINNGGYT"

BASE COUNT 303 a 228 c 341 g 383 t

ORIGIN

```
1 atgccgccga tgggtacgct agagcacgtg gtgtgggtgc gtgtgttggt acgtttactg
61 ttgggggatt aagtgttttg aatgctatag cgggtgctta tagtgagaat ttaccgttga
121 tatgtattgt tgggggacct aattcgaatg attatggaac caacaggatt ttacatcata
181 ctattggggt gcaggatttt agtcaggagc ctggttgctt tcaaactgtc acttgttacc
241 gggctgtggt gaataattta gaggatgcac atgaactgat tgacactgct gtatctacag
301 ctttgaaaga gagtaagccg gtgtatataa gcataggctg taatctgccg gggattccac
361 accctacttt cagtctgtaa ccagttccat ttgccctctc tcccagactg agtaatatga
421 tgggtttgga agcagcagtg gaggcagcgg ctgagttctt gaacaaagca gtaaagccag
481 tacttgttgg agggccaaaa atgcgggttg caaaggcatc tgatgctttt gttgagttgt
541 ctgatgcctg tggttatgct gtgcagtgta tgccatctgc taaaggggtg tttcccgaac
601 accattcaca tttcattggg acttactggg gtgcagtgag cacagccttc tgtgctgaaa
661 tcgtggaatc tgctgatgca tacttgtttg ctggacctat ttttaatgac tatagctctg
721 ttggttattc tctgcttctc aagaaagaga aagcgatcat tgtccagcct gatcgtgtga
781 ctattgggaa tggaccagcg tttggttgtg ttctgatgag ggatttcctt gctgcattag
841 caaagagatt aaagcataat ccgactgctt ttgagaatta ccataggatt tatgttctctg
901 agggacatcc tcttaaagt gaggctaagg aggcattaag ggttaatgtt ctatttcaac
961 acattcagaa tatgttgtct ggcgacagtg ttgtgattgc tgagacgggg gactcgtggt
1021 tcaattgcc aagctgaaa ttgccaaaag gatgtgggta tgagttccaa atgcagtatg
1081 gatecatcgg ttggtctgtc ggtgcaactc ttggttacgc acaagctgca cctgaaaagc
1141 gggtgattgc ttgcatcggg gacggtagct ttcaggtgac tgctcaggat atttcaacga
1201 tgctgaggtg tggacaaaga accattatct tcttgataaa caatggtggc tatac
```

//


Revised: July 5, 2002.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Oct 3 2002 17:48:23



**Nucleot**



PubMed

Nucleotide

Protein

Genome

## Structure

PonSet

### Taxonomy

Search **Nucleotide**  for

**Nucleotide**

for

\_\_\_\_\_

Go

Clear

## Limits

[Preview/Index](#)

## History

## Clipboard

## Details

## Display

**default**

# Save

Text

Add to Clipboard

## GetSubsequence

□ 1: U71122. *Arabidopsis pyruv...*[gi:1616786]

## Links

```

LOCUS       ATU71122                4069 bp    DNA        linear    PLN 08-OCT-1996
DEFINITION  Arabidopsis pyruvate decarboxylase-2 (Pdc2) gene, complete cds.
ACCESSION   U71122
VERSION     U71122.1   GI:1616786
KEYWORDS    .
SOURCE      Arabidopsis thaliana.
  ORGANISM  Arabidopsis thaliana
            Eukaryota; Viridiplantae; Streptophyta; Embryophyta; Tracheophyta;
            Spermatophyta; Magnoliophyta; eudicotyledons; core eudicots;
            Rosidae; eurosids II; Brassicales; Brassicaceae; Arabidopsis.
REFERENCE   1  (bases 1 to 4069)
  AUTHORS   Dolferus,R., Peacock,W.J. and Dennis,E.S.
  TITLE     Two pyruvate decarboxylase genes from Arabidopsis
  JOURNAL   Unpublished
REFERENCE   2  (bases 1 to 4069)
  AUTHORS   Dolferus,R., Peacock,W.J. and Dennis,E.S.
  TITLE     Direct Submission
  JOURNAL   Submitted (18-SEP-1996) Plant Industry, C.S.I.R.O., G.P.O. Box
            1600, Canberra, Act. 2601, Australia
FEATURES             Location/Qualifiers
     source            1..4069
                       /organism="Arabidopsis thaliana"
                       /strain="Landsberg-erecta"
                       /db_xref="taxon:3702"
                       /chromosome="5"
                       /map="between DFR (7.5 cM) and RFLP m435 (1.8 cM)"
     gene            1447..3270
                       /gene="Pdc2"
     CDS             1447..3270
                       /gene="Pdc2"
                       /EC_number="4.1.1.1."
                       /codon_start=1
                       /product="pyruvate decarboxylase"
                       /protein_id="AAB16855.1"
                       /db_xref="GI:1616787"
                       /translation="MDTKIGSIDACNPTNHDIGGPPNGGVSTVQNTSPLHSTTVSPCD
ATLGRLYLARRLVEIGVTDVFSVPGDFNLTLDDHLIAEPNLKLIGCCNELNAGYAADGY
ARSRGVGACVVTFTVGGLSVLNAIAGAYSENPLICIVGGPNSNDYGTNRILHHTIGL
PDFTQELRCFQAVTCFQAVINNLEEAEHELIDTAISTALKESKPVYISISCNLPAIPLP
TFSRHPVPFMLPMKVSNIQGLDAAVEAAAEFLNAVKPVLVGGPKMRVAKAADAFVEL
ADASGYGLAVMPSAKGQVPEHHKHFIGTYWGAVSTAFCAEIVESADAYLFAGPIFNDY
SSVGYSLLLLKKEKAIIVQPDRVITIGNPAFGCVFMKDFLSELAKRIKHNNTSYENYHR
IYVPEGKPLRDNPNESLRVNVLFQHIQNMLSSESAVALAETGDSWFNCQKLKLPEGCGY
EFQMQYGSIGWSVGATLGYAQAMPNRRVIACIGDGSFQVTAQDVSPMIRCCHKTIIFL
INNNGYTI EVEIHDGPYNVIKNWNYTAFVEAIHNGEQKWTPKVRCEEELVKAINATAT
NEEKESFCFIEIVHKDDTSKELLEWGSRSVSAANSRPPNPQ"
BASE COUNT   1105 a      828 c      855 g      1281 t

```

## ORIGIN

```
1  aagcttttaa ctttttccgt caatttcggt gtaaaacaca gagacataga agacagagag
61  aaaaaacaca gacctgcgaa tttttcgtct ccttttcttc agctttcttc ttgtggcatt
121 taacaacgag tcaataaagg gctaatacga aaaaagacaa aaacatttga gagggccaaa
181 aatattttta ttacagcaaa aaaaattgtg gatttgtttg gttataacac tttctctgtt
241 ttgttaaatt aatggtttgt ttttgtttgg gcttttagat aatatttttg cttaaagata
301 tttccttctt gcttttttat tatgcttctg tgatcagtta cttcttcttt ttttagaggc
361 acgtgaaaac cacgtgtcta ctatgcaaag atctttgatt taccaggcca ttcattagct
421 aagcaatatt ataatacatc ttcattcttt ggattagccc ggagataact actttaaaag
481 cccattcaat catggactta aacttagtat tgggtatatc acaaatttcg gttcaacttc
541 attcagatta gttgctaaat tattgtttat ctgttctctg atacgttcgt tactttcata
601 cattttcaac catacaataa accaaatcat atttcagatt tattccaaat atagtggagc
661 acattcaatt tcccaaagta catttgattg aatgctcgag cttcttatct tatcccttga
721 tccatcaaat tatcagcttc ttctaaccat tctatttctg ttcacaatct tactttcttc
781 ttttgcaagt aattaaagga agagcttgag gaattaggac aatcgaaata tttcttatca
841 ttgcaaaagt caataacaaa accaaagctt ggtgaaagag attgaagaat ctgatgatcg
901 agagagcgag aaagagattg tgggagatat gtgattgaag attggggatc tacggatttt
961 tgtaatcgag tgagagcggc gggaaaatcg gcgttagggg ggaaatacga gtctctttag
1021 ggtttcgaaa agtcgccaag agcgggagtt tctttttgag tcacgtggta ttgacacgaa
1081 ccgggtcgga tcctatatcc aatcgggtac tatgttgccc ttgtgggcta cgcgtgtcgc
1141 gtgagagcaa ctatctctgc ctactcacia ctattctcaa cgtgaacttt ctaacttggt
1201 ctccaatatc caactatata aggacacgtg tctaattctc gagacacata aatatctgcg
1261 gctaaaaata tcttatttag cgctattttc ataccaccgt ttgattactt tagccaccgt
1321 tgttttacgt ctataattaa gtatcttcca ctaatctcat attttcaaac tgtactttta
1381 cctcgagctc tctccaattt tctcatagct ctattgttcg aaaattctga aacgaacggt
1441 accactatgg aactaagat cggatctatc gacgcgtgta acccgacca ccacgatatc
1501 ggcggtcctc caaacggcgg agtctccacc gttcaaaaca caagtcact tctactccac
1561 accgtcagcc cctgcgacgc gactcttggc cgttacctag caagacgggt agtcgaaatc
1621 ggcgtcaccg atgtcttctc cgttcctggg gatttcaacc tgacgcttct cgatcaccta
1681 atcgccgaac caaacctcaa gctgatcggt tgctgcaacg agcttaacgc cggatacgtc
1741 gctgacgggt acgctagatc tcgcgggtgt ggtgcgtgcg tcgttacgtt caccgtcggt
1801 ggattgagtg ttctgaatgc gatcgccggg gcttacagtg agaactctgc cctgatttgc
1861 atcgctcggt gtccaaactc caacgattac ggtaccaata ggattcttca tcatacaatt
1921 ggtttacctg atttcaactc agagcttagg tgttttcaag ctgttacttg ttttcaagct
1981 gtgattaata acttagaaga ggctcatgaa ctatcgata ctgcgatttc aactgctttg
2041 aaagaaagca aacctgttta tatcagtatc agctgtaatt taccggcgat tcctcttccg
2101 acgttttagt gtcacacctg tccgttcatg cttccgatga aggttagcaa tcagattggg
2161 ttagatgcgg cgggtggagg agctgctgag ttcttgaaca aagctgtgaa gccagttctt
2221 gttggtgggc cgaaaatgcg ggttgcgaaa gccgcggatg cttttgttga gcttgctgat
2281 gcttctggct atggtcttgc tgtgatgcct tctgctaaag gacaagtacc tgagcatcac
2341 aagcatttta tagggacgta ttggggagct gtgagtacag ctttttgtgc tgaaatcggt
2401 gaactctcgg atgcttatct gtttgcaggc ccgattttca acgattacag ttctgttggg
2461 tattctctgc ttctcaagaa ggagaaggca atcatcggtc agcctgatcg ggttactatc
2521 ggtaacggac ctgcgtttgg atgtgttttt atgaaggatt ttctaagcga gttggctaaa
2581 cgaattaaag acaacaacac ttcttatgag aattatcaca ggatctatgt cccagaagga
2641 aagcctttga gagataaccc gaatgagtct ttgaggggta atgtactgtt ccaacacatt
2701 cagaatatgc tctcttctga gtctgctgtg cttgctgaga caggagattc ctgggttcaac
2761 tgtcagaagc tgaagctccc tgaaggatgc ggttacgaat tccaaatgca gtacggatca
2821 attggctggg cagtgggtgc tactctaggc tatgctcaag ccatgccaaa caggcgtgtc
2881 attgcttgta ttggagatgg tagtttccag gtaaccgcac aggatgtatc tccgatgata
2941 cgggtgtggg acaagaccat aatcttctc atcaacaacg gaggctacac cattgagggtg
3001 gaaattcacg atggtcctta caatgtcata aagaactgga actacacagc ttttgttgag
3061 gccatacaca atggagaagg aaaatgctgg actcccaagg tgagatgcga ggaggagtta
3121 gtgaaagcaa tcaacacggc aaccaatgag gaaaaagaga gcttttgttt cattgaagtg
3181 atagtgcaca aagacgatac aagcaaggaa cttttggagt ggggctctag agtctctgct
3241 gctaatagtc gtcccccaaa tccgcagtag agtatataaa tgcactcaac ttatatatat
3301 ttcagatttg gtagtgtctt caccgttcta tgtaaagtag tgtgtatcct tttacacca
3361 tctggatgga aaaaaaatgt gtgtccccct gaggatgtaa actgttttag tgccaaata
3421 agaacgtgtt tcctctgctt catcttctgt cctcagtttt acgcagattt atcagagcta
3481 tagtctttgt gtcttggatt ggtatgttga ttgacttaag aaagttgaga tccatcccaa
```

```
3541 gatgccaaat gattacttta aaggcagttg gtagagtcct ttcttttccc aatttatact
3601 aacttttggt ttaccttttc ttgctttctg tttttttggt tcatggctaa atttgatctt
3661 gtttctatga ggaatgcagt gaggcgtctt gggagtgatc cactggttta caaaatactt
3721 catcgctatg ctggttgaat gttacgaatc tttcgctggt ggccataatt ttattactta
3781 agtaggatta aatagcttat tccatagtaa ctctcctaaa tttctgcata gagaaacctg
3841 tccaaatttg tgatgcagct tgaattcgac aggaattggg gttaatgcct caagtcggt
3901 gaagatgcag aaaccttggt actaactggc catgaactct acacaatcga cgtacctagt
3961 aatatcaatg agataaaacc aggccagtac ataactgtga ctactactga cactgctaaa
4021 tcttttgtct gcactttgct gttagacaca tagttaatca caaaagctt
```




//

Revised: July 5, 2002.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Oct 3 2002 17:48:23



[PubMed](#)
[Nucleotide](#)
[Protein](#)
[Genome](#)
[Structure](#)
[PopSet](#)
[Taxonomy](#)

Search  for

☐ 1: U65927. *Neurospora crassa*...[gi:1655908]

[Links](#)

LOCUS NCU65927 3741 bp DNA linear PLN 01-NOV-1996  
 DEFINITION *Neurospora crassa* pyruvate decarboxylase (cfp) gene, promoter region, and complete cds.  
 ACCESSION U65927  
 VERSION U65927.1 GI:1655908  
 KEYWORDS  
 SOURCE *Neurospora crassa*.  
 ORGANISM *Neurospora crassa*  
 Eukaryota; Fungi; Ascomycota; Pezizomycotina; Sordariomycetes; Sordariales; Sordariaceae; *Neurospora*.  
 REFERENCE 1 (bases 1 to 1699)  
 AUTHORS Temporini, E., Alvarez, M., Mautino, M., Kawaguchi, T., Kinghorn, J. and Rosa, A.  
 TITLE Cloning, sequence and functional analysis of the *Neurospora crassa* pyruvate decarboxylase gene promoter  
 JOURNAL Unpublished  
 REFERENCE 2 (bases 1 to 3741)  
 AUTHORS Temporini, E.  
 TITLE Direct Submission  
 JOURNAL Submitted (02-AUG-1996) Esteban Temporini, Facultad de Ciencias Quimicas-Univ. Nac. Cordoba, Departamento de Quimica Biologica, Ciudad Universitaria s/n, Cordoba 5016, Argentina  
 FEATURES  
 source Location/Qualifiers  
 1..3741  
 /organism="Neurospora crassa"  
 /strain="wild type"  
 /db\_xref="taxon:5141"  
 /chromosome="VIIR"  
 /clone="x10c4"  
 /tissue\_type="mycelial"  
 /clone\_lib="pMOCosX of Marc Orbach"  
 protein\_bind 456..460  
 /note="This is a sequence with high homology to the binding site for GCR1 protein of *Saccharomyces cerevisiae*"  
 /bound\_moiety="GCR1 protein"  
 /function="transcriptional activation of glycolytic genes"  
 protein\_bind 645..649  
 /bound\_moiety="GCR1"  
 /function="transcriptional activation of glycolytic genes"  
 protein\_bind 741..753  
 /note="sequence with high homology to the binding site of RAP1 protein of *Saccharomyces cerevisiae*"  
 /bound\_moiety="RAP1"  
 /function="transcriptional regulation"  
 protein\_bind 1003..1015  
 /bound\_moiety="RAP1"  
 /function="transcriptional regulation"

protein\_bind 1370..1382  
 /bound\_moiety="RAP1"  
 /function="transcriptional regulation"  
protein\_bind 1384..1396  
 /bound\_moiety="RAP1"  
 /function="transcriptional regulation"  
protein\_bind 1390..1394  
 /bound\_moiety="GCR1"  
 /function="transcriptional activation of glycolytic genes"  
misc\_signal 1602..1609  
 /function="transcription initiation signal in *Neurospora crassa*"  
gene 1697..3409  
 /gene="cfp"  
CDS 1697..3409  
 /gene="cfp"  
 /codon\_start=1  
 /evidence=experimental  
 /product="pyruvate decarboxylase"  
 /protein\_id="AAB17969.1"  
 /db\_xref="GI:1655909"  
 /translation="MVAQQQGKFTVGDYLAERLAQVGVRRHFFVVPGDYNLILLDKLQA  
 HPDLKEVGCANELNCSLAAEGYARANGISACVVITYSVGALSAFNGTGSAYAENLPLVL  
 ISGSPNTNDPSQYHILHHTLGHPDYTYQYEMAKKITCCAVAI PRAIDAPRLIDRALRA  
 AILARKPCYIEIPTNLGATCVRPGPISAITDPITSDKSALEAAAKCAA EYLDGKLP  
 VILVGPKAGRAGSEKELIEFAEAMGCAVALQPAKGMFPEDHKQFVGIFWGVSSDAA  
 DAMVHWADAMICVGAVFNDYSTVGWTA VPNIPLMTVDMDHVTFPGA HFSRVRMCEFLS  
 HLATQVTFNDSTMIEYKRLKPDPPHVHTAERE EPLSRKEISRQVQEMLTDKTSLFVDT  
 GDSWFNGIQLKLPPGAKFEIEMQWGHIGWSIPAAFGYALRHPDRHTIVLVGDGSFQVT  
 AQEVSQMVRFKVPITIMLINNRGYTIEVEIHDGSYNKIKNWDYAMLVEAFNSTDGHAK  
 GLLANTAGELADAIKVAESHKEGPTLIECTIDQDDCSKELITWGHYVAAANARPPRNM  
 SVQE"

BASE COUNT      845 a    1139 c      943 g      814 t  
 ORIGIN

```

1  ccgcggcctc ccactcagaa taacagtgcg cgtcatggta gaacaagcat atggggagca
61  ggccttttgg caacaaaggt ggtcaatctc atggtcgtca ttctgtcaca tctcgcggtt
121 gtgggaggag gttgagcaga cagaaaagta aatgggtaca cgaggggtgg tgggtgcctt
181 ttgttggtcg ggcgccagg cttttccacc cctctctcca tagcagatat ggcacatgtg
241 tttaactcaa taggtcatt tggtactata aatacaacct tacgccaaac ctttcatcga
301 cgaaacttgt accaggaata tctatatctt cattgacttt tcaaggacgg aggttccaat
361 gatcagctct cgtcctaagc tcaaatgata ttggttggtat tctatttggg aaagaagcag
421 ccgccctcat ctccgttgcc ttggttgagg cggcacttcc gagggacaga ggtgaaagtg
481 ttaggtcgca gtagggcgaa tctcatttcc caagtcagac aactttgaca tgatacgatg
541 tgcaaactct tttcttctga atctctgctc ttatcttaag tttcttggaa tttcaaatac
601 acaagagtgg ttgaatggag cttggtgggc tgttggttaa caggcttcca gcactgggt
661 tgcgtttctg actatcgccc tcaataaaaa catgttccat attttctatc ctcatgcctt
721 tggcactcag actcaagagc gcacccacca ccctcagggg acctaggcat agcttcagta
781 tcgaactcat gtcaacgcgg agcattgacc acatcgacct caaacctcaa caaacagAAC
841 aaccgacatg agctcagccc acgttgccat tcggcagacg gagccgaatc gtgctgactg
901 ttttgaaaat ttaccgcccg atgggtgccg ctcggaaggc cagcctgagg catctgtgtc
961 tggcatgtcg ccgacgcggg taaggtccgc ttttctgggg cgcaacccca ccagccgcca
1021 atgggaaggg ccggtgtggc ttgggtgcga gtgcctggat tgcggcgggt gtctgatcaa
1081 atcgtcgaat gctcgctgca tatcggaacc aacgaggccg cttatctttc cctacaacac
1141 ttcactctga ccgacgtcga acaaccggag ccagcagcgg gatggccctg tgcaacgtcg
1201 taatccatgg tccccactc ctgtctatga tgcattgccc gcatgctctg acttcacaac
1261 ctcgtaacag aatgttcaca ccaccagcgg accgtgggtg cgcttggtct gatggccctg
1321 tcaagcaact ggcgaccgtc tggccattgc tcccacgaa cgacaccatg cagccatgca
1381 tcagcacacc ttccttggtc cttggcccat gcgagagcgc cggagcgtct tcggggacat
1441 gtcacgcggg agatggcctc aagccatcga tggcaatcct gtagcaggga ggccatcggg
1501 acaagctctc tgagccaatc atcacatcac cgtggtcaac acaggacaat aaaaggcttc
  
```

```
1561 tcgcagcccg ccgctcctctt gggtagccga cttctgcatg gtcatgaaag tacagcttcg
1621 aactaccac tactactacc acctccagtc tttgacactt cttcaccacc agatctcacc
1681 cttaaagagag agcatcatgg tagcccaaca acaaggaaag ttcacggtgg gcgactacct
1741 cgccgagcgt cttgtctcagg tcggcgctcg ccaccacttt gtcgtccccg gcgactacaa
1801 cctcctcttc ctcgacaagc tacaagctca cccggatctc aaagaggttg gttgcgccaa
1861 cgagctcaac tgcctctctt cgccgaagg ccatgccctg gccaacggga tctccgctg
1921 cgttgtcact tacagcggtt gtgccctctc ggccttcaac ggcaccggca gtgcctacgc
1981 agagaacctc cccttgggtc tcatcagcgg ctgcaccaac accaacgacc ccagtcaata
2041 tcacatcctt caccataccc tcggccaccc ggactacacc taccagtatg agatggccaa
2101 gaagattacc tgcgtgcgtg tggccattcc ccgcgccatc gatgcacccc gcctgatcga
2161 ccgtgccctc cgcgcgcgca tcttgcctcg gaagccctgc tacatcgaga tccccaccaa
2221 cctcgccgga gctacctgcg tgcgccccgg ccccatcagc gccatcaccc accccatcac
2281 cagcgacaag agcgcctctt aggcgcgcgc caagtgcgct gccgagtatc tcgacggcaa
2341 gctcaagccc gtcactcttg ttggcccaaa agctggccgt gccggttcgg agaaagagct
2401 catcgagttc gccgaggcca tgggctgtgc tgttgccctg caacctgccg ccaagggcat
2461 gttccccgaa gaccacaagc agttcgtcgg catcttcttg ggacaagtta gctccgatgc
2521 cgcagacgcc atggtgcaat gggcgacgc catgatttgc gtcggcgccg tctttaacga
2581 ctacagcacg gtcggatgga ctgccgtgcc caacatcccg ctcatgaccg tcgatatgga
2641 tcatgtcacc ttccccggcg cacacttcag ccgcgtccgg atgtgcgaat tctcagcca
2701 cctggccacc caggtcacct tcaacgactc caccatgac gagtacaagc gcctcaagcc
2761 cgaccccccg cagtgacaca cggcggaacg cgaggagccg ctctcgcgca aggagatctc
2821 gcgacaggtg caggagatgt tgacagacaa gaccagcctg ttcgtcgaca cgggcgactc
2881 gtggttcaac ggcattccagc tgaagcttcc tcttggcgcc aagttcgaga tcgagatgca
2941 atggggccac atcggtcgtt cgatccctgc ggcctttggc tacgccttgc gccacctga
3001 caggcacacc attgtgcttg tcggcgacgg atctttccag gtgacggccc aggaggtgtc
3061 gcagatggtg cgcttcaagg taccatcac catcatgtc atcaacaacc gcggtacac
3121 gatcgaggtc gagatccacg acggttctta caacaagatc aagaactggg actacgcat
3181 gttggtggag gccttcaaca gcaccgacgg ccacgcaaag ggcctgctgg ccaacacggc
3241 cggcgagctc gctgacgcta tcaaggtggc tgagagccac aaggagggtc ctacgtgat
3301 cgagtgcacg attgatcagg atgactgcag taaggagctc atcacctggg gtcattacgt
3361 ggctgctgcg aatgcgaggc ctccccgcaa catgtcgggtg caggagtaag gaagatgcta
3421 atgtggatag gcatgttgaa atgacggatg tgataccaaa tgatagtatt atcaaagata
3481 caacggcggc ggcggcggcc gaggtggtgg tagacaatgg gccacgatat agtgtatgga
3541 aaaaggtcaa acggggcttc aagtttttgt catgtcaata aatgttttgt tttcctttta
3601 ggctttctca tagatcgttc atctcatctc aatttctata tatctctctc tctcattca
3661 tgtctctgtc ccaccgggtt tggttaaaat ctatacaaag acaagtggta aacgaacaag
3721 aaaagctatt caatcgaatt c
```

//

Revised: July 5, 2002.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Oct 3 2002 17:48:23



```
241 caggaagtgt acacgatggt gcggtgcggg cagaagagca tcattcttct catcaataat
301 ggtgggtaca ccatcgaggt ggagatccac gacggggcgt acaacgtcat caagaactgg
361 gactacaccg gcctcatcga cgcaatccac aactctgacg gcaactgctg gacaaagaag
421 gtccggactg aggaggagct gatagaggcg atcgcgacgg caacggggcg caagaaagat
481 tgcctctgct tcatcgagat catcgtgcac aaagatgaca cgagcaaaga gcttctcgag
541 tggggatcca ggggtatcggc tgccaacagc aggcgcgcga atccccagtg attttgctag
601 ctggtcttct catgacgcac actcgaatgc gctagtctgc caatgtccga gatatgctgc
661 ccattatcgt ctgtgaatta attgtctaata gcgtctgcat ccagttaaat tttctcagcc
721 gagaggttga agtatttatg tagatggggg tgcccgctct atgattatgt ttgtaattca
781 atatgcttgg cattaatggt ttgtaaacct gtaatgtatt gcgatgaaat aacatggacg
841 atttgaagga gaagaa
```

//

Revised: July 5, 2002.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Oct 3 2002 17:48:23



```

YRRIFVPDREPPNGKPNELRVNVLFKHIKGMLSGDSAVVAETGDSWFNCQKLRLPEG
CGYEFQMQYGSIGWSVGATLGYAQAADKRVIAICIGDSFQVTAQDVSTMLRCGQKSI
IFLINNGGYTIEVEIHDPYNVIKNWDYTGLVNAIHNSGNCWTMKVRTEEQLKEAIA
TVTGAKKDCLCFIEVIVHKDDTSKELLEWGSRVSAANSRPPNPQ"
exon          <1415..1954
               /gene="Pdc gene"
               /number=1
intron        1955..2044
               /gene="Pdc gene"
               /number=1
exon          2045..2977
               /gene="Pdc gene"
               /number=2
intron        2978..3083
               /gene="Pdc gene"
               /number=2
exon          3084..3272
               /gene="Pdc gene"
               /number=3
intron        3273..3366
               /gene="Pdc gene"
               /number=3
exon          3367..>3537
               /gene="Pdc gene"
               /number=4
repeat_region 3579..3596
               /note="tgcgtc repeat"
               /rpt_type=TANDEM
polyA_signal  3790..3795
polyA_signal  3814..3819
BASE COUNT    898 a   1197 c   1003 g   942 t
ORIGIN
1  ctcaactgtcc atgccatcag atgggtactgt acaggaagaa tgatactgta gttaagtacc
61  cccaaaatat aaggcatttt caagattatt gaaaagtcaa agtggtgttt caagtttgac
121 caaatattata taacgaagta ataacgttta cgatattaaa taaatgtata ccattagatt
181 ctctattataa tatattttta tactatatct actctgtgtc ataaatataa atatttatga
241 ttctcgttat aatttttaatc aaacttttaa aactttgact gtctaaattt ttgtaaatga
301 cttatatattt gggacagagg agagtaaggt tttagcctcg gcggcactgg aagaatgtta
361 attagaagca tgcccacgtc atctttttgt tactgtattt atctaagtga atatcatgta
421 ataaataaac acatatagct atatatattt ttgttaattt taagtgcaga ccgaatagta
481 gtttataaag agtaaagtac actaccgacc taaaacttg gcatgcgggt aaacacgtac
541 tgcgaactttg tcagccaata aaaaaccatc caaaaccgag ttgtgaaaac cagatgccga
601 cgtgggtcaga gccacttaag atgtctaact tggtttatat gttacaaagt tacatcttaa
661 tttatcttta gcttccaaca tcctatctac atgtgtctag ctagagttac tcaccacact
721 cgatcgggaa catggctaga gttgtcactg tatttctcta atgataatgt catgtaattt
781 ttttctgtga ctgtcagtgg tcaccgaggt ttaagggagc aacttaattt aggacgggat
841 ttttaataac ggtataacat ttttggcacg cacaagcaca aaaccagtac cgccgcagtt
901 tgttttggtg caactgctga tactagcagc tacagtgaat cctatctgta gtgtgctggg
961 gctatatgta gctgtagtgt gcaacggtaa tactaggaca tggaaacctg gaggcggcca
1021 gctaggctag tactgtacta ctaggacaac agctcagctc aggcaacgcg ccggaaggca
1081 cacgaagccc aagtttggtt tgggtggctcg tttatcccag cccacagcat gcgaacagac
1141 tggcccgctt tgccttggcg gcttggcact gtgtgcagtg taacaacctt tcgcccttcc
1201 cctccctcag ggcttcgggc cttctgccct tctgccttca gcttcagacc cgttcctgtc
1261 tatataaatc cccacgatgg gaggggttgc ccacccacc catcatcagt cagtcagtca
1321 gcagccgcaa cctcgtacaa tcacgaaagc agcacacgag tccacaacag caccctcctt
1381 ctttttcttc acgtgcgttg cgtgccctgc ccccatggag accctccttg ctggcaacct
1441 cgccaacggc gtggcggaagc cgacgtgcaa cggcgtcggc gccctgcccc tggccaactc
1501 ccacgccatc atcgccacgc cggcgggcgc tgcggcgacg ctggccccgg ctggcgccac
1561 gctcggccgg cacctggcgc gccgcctcgt gcagatcggc gcctccgacg tcttcgccgt
1621 cccggggggac ttcaacctca ccctgctcga ctacctcatc gccgagccgg gctgaccct

```

1681 cgtcgggtgc tgcaacgaac tcaacgcggg gtacgcgcgc gacgggtaac cgcgggtcgc  
1741 gggcggtcggc gcctgcgcgc tcacgttcac cgtcgggtggg ctacagcgtgc tcaacgccat  
1801 cgcgggcgcgc tacagcgaga acctccccgt cgtctgcate gtgggcgggc ccaactccaa  
1861 cgactacggc accaaccgca tccctcacca caccatcggc ctccctgact tctcccagga  
1921 gctccgctgc tccagacga tcacctgcta ccagggtgctg cgcgctaata ctgctgtgtg  
1981 cttctcatct tctctcttct gctagctgct cggcatcgct atatcttctc tctactctcg  
2041 gcaggccatc atcaacaacc tggacgacgc gcatgagcag atcgacacgg ccacgcgcac  
2101 ggcgctgagg gagagcaagc ccgtgtacat cagcgtcagc tgcaacctgg ccggcctctc  
2161 ccaccgcacc ttcagccggg atccgggtgcc catgttcate tcgcgcaggc tgagcaacaa  
2221 ggccaacctg gtagtacgcg tggaggcggc ggccgacttc ctcaacaagg cgggtgaagc  
2281 ggtgatggtg ggcgggcccga agatccgggt ggccaaggcc agggaggcgt tcgcgcgcgt  
2341 cgcggacgcc agcgggtacc ctttcgcgt catgccggcc gccaaaggcc tgggtgcgga  
2401 gcaccaccgc cgtttcatcg gcacctactg gggcgccgctc agcaccacct tctgcgcgga  
2461 gatcgtggag tccgcgcacg cctacctctt cgcggggccc atcttcaacg actacagctc  
2521 cgtcgggtac tccctcctgc tcaagcggga gaaggccgctc atcgtgcagc ccgaccgcac  
2581 ggtggtcggc gacggcccg cgttcggctg catcctcatg cccgagttcc tccgcgcgct  
2641 cgccaagcgc ctacggcgca acaccacggc gtacgacaac taccgcgcga tcttcgtccc  
2701 cgaccgcgag ccgcccacg gcaagcccaa cgagccgctt agggtaacg tcctcttcaa  
2761 gcacatcaag ggcattgctgt ccggcgactc cgcggctcgtc gccgagaccg ggcactcgtg  
2821 gttcaactgc cagaagctca ggctccccga gggtgcggc tacgagttcc agatgcagta  
2881 cggctccatc ggggtggtcgg tcggcgcgac gctcgggtac gccaggcgg ccaaggacaa  
2941 gcgtgtcatc gcgtgcacg gcgacggaag cttccaggta caatacggca catccatcca  
3001 tggcacacat gctgtctctc tgtctgtgct tgtacgtgct cactgctgat cggacgatcg  
3061 acgacgtgtg gctgggtgcc cagggtgacg cgcaggacgt gtccacgatg ctccgggtgcg  
3121 ggcagaagag catcatcttc ctcatcaaca acggcggtta caccatcgag gtggagatcc  
3181 acgacggccc gtacaacgtg atcaagaact gggactacac gggcttggtc aacgccatcc  
3241 ataactccga gggcaactgc tggaccatga aggtctgtcg atcagacaga cctctctcca  
3301 tccaacactg aaccccacca ccaccagctg cgcgctaata aaaccgctcc accgtgtgcc  
3361 ctgcagggttc ggacggaaga gcagctcaag gaagcgatcg ccacgggtac gggcgccaag  
3421 aaagactgcc tctgcttcat cgaggtaatc gtgcacaagg acgacacgag caaggagctc  
3481 ctcgagtggg ggtccagggt ctccgcgcgc aacagcaggc cgcaccaacc ccagtgatga  
3541 gcccgtcggg cgggtctcgag cgtttctgca atggcgctg cgtctgcgct tgcgtctgtc  
3601 tgccctgaaa atttttcggg tcgtttactc gttttcagtc agttactcag tttgtccgctc  
3661 ttcgtcgagc agaactacat cctgctgag ggaagctacc aatctttatc ctacgatgat  
3721 aatgtttgca atttcacttg ctcccgattg agttctgtta taaatctttg caacatcttt  
3781 tctcttatca ataaaaattt tttttctctt atcaataaaa tttgttcgcg acgcccgtct  
3841 ccttagagac ctcggttact gtgcgcggg tgtgtatccc catgacctag acggacttaa  
3901 gcctgaatcg tgtggaatag gtgatttagg gttagtttag caacccttt ttttaagagat  
3961 ttctatcttt taagatagat tagtttattt tcctttgagg aaaacgaaat tccgtaagaa  
4021 attatagtat taaactagcc


//

Revised: July 5, 2002.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Oct 3 2002 17:48:23





[PubMed](#)
[Nucleotide](#)
[Protein](#)
[Genome](#)
[Structure](#)
[PopSet](#)
[Taxonomy](#)

**EXHIBIT**

H

Search Nucleotide for 
Go
Clear

Limits
Preview/Index
History
Clipboard
Details

Display
default
▼

Save
▼

Text
▼

Add to Clipboard

Get Subsequence

**□ 1: U75311. Pichia stipitis p...[gi:2734882]**

## Links

```

LOCUS       PSU75311                2305 bp    DNA         linear     PLN 19-FEB-1998
DEFINITION  Pichia stipitis pyruvate decarboxylase 2 (PsPDC2) gene, complete
            cds.
ACCESSION   U75311
VERSION     U75311.1   GI:2734882
KEYWORDS    .
SOURCE      Pichia stipitis.
  ORGANISM  Pichia stipitis
            Eukaryota; Fungi; Ascomycota; Saccharomycotina; Saccharomycetes;
            Saccharomycetales; Saccharomycetaceae; Pichia.
REFERENCE   1  (bases 1 to 2305)
  AUTHORS   Lu,P., Davis,B.P. and Jeffries,T.W.
  TITLE     Cloning and characterization of two pyruvate decarboxylase genes
            from Pichia stipitis CBS 6054
  JOURNAL   Appl. Environ. Microbiol. 64 (1), 94-97 (1998)
  MEDLINE   98096791
  PUBMED    9435065
REFERENCE   2  (bases 1 to 2305)
  AUTHORS   Lu,P., Davis,B.P. and Jeffries,T.W.
  TITLE     Direct Submission
  JOURNAL   Submitted (16-OCT-1996) Institute for Microbial and Biochemical
            Technology, USDA, FS, Forest Products Laboratory, One Gifford
            Pinchot Drive, Madison, WI 53705, USA
FEATURES             Location/Qualifiers
     source            1..2305
                     /organism="Pichia stipitis"
                     /strain="CBS 6054"
                     /db_xref="taxon:4924"
     TATA_signal       216..222
     TATA_signal       267..273
     gene              393..2102
                     /gene="PsPDC2"
     CDS                393..2102
                     /gene="PsPDC2"
                     /EC_number="4.1.1.1"
                     /function="decarboxylation of pyruvate during ethanol
                     fermentation"
                     /note="thiamine binding protein"
                     /codon_start=1
                     /product="pyruvate decarboxylase 2"
                     /protein_id="AAC03165.1"
                     /db_xref="GI:2734883"
                     /translation="MVSTYPESEVTLGRYLFERLHQLKVDITFGLPGDFNLSLLDKVY
                     EVPDMRWAGNANELNAAAYAADGYSRIKGLSCLVTTFGVGELSALNGVGGAYAEHVGLL
                     HVVGPVSISSQAKQLLLHHTLGNLDFTVFHRMSNSISQDTAFSLSDISIAPGQIDRCIR
                     EAYVGHQRPVYVGLPANMVDLKVPSSSLLETPIDLKLKQNDPEAQEVVETVLKLVSQATN
                     PIILVDACALRHNCKEEVKOLVDATNFOVFTTPMGKSGISESHPRLGGVYVGTMSSTP

```

VKKAVENADLILSVGSLLSDFNTGSFSYSYKTKNVVEFHSDYMKIRQATFPGVQMKEA  
LQOLIKRVSSYINPSYIPTRVPKRKQPLKAPSEAPLTQEYLWSKVSGWFREGDIIVTE  
TGTSAFGIIQSHFPSNTIGISQVLWGSIGFTVGATVGAAMAAQEIDPSRRVILFVGDG  
SLQLTVQEISTLCKWDCNNTYLYVLNNDGYTIERLIHGKSASYNDIQPNHLSSLRLF  
NAKKYQNVRVSTAGELDSLFSKKFASPDRI RMIEVMLSRLDAPANLVAQAKLSERVN  
LEN"

misc\_feature

1734..1799

/gene="PsPDC2"

/note="encodes thiamine binding region"

BASE COUNT        681 a        472 c        460 g        692 t  
ORIGIN


```
1  aagaattgct  tcttaccttg  atggcattcc  taatgaaacc  atccactcaa  taggctgtgt
61  tgaatgctgc  ctaatcgcat  atcctacaca  taataggaat  gtctccagct  tatggacgat
121 taaggaacca  ttcattgcgtt  taaacatgct  gcacattaat  cttttctaca  agttcatcac
181 gaactgatgg  gaaatgcctt  cagatgccca  gggcctataa  tcgacaactg  tactagecaa
241 taagcagctg  atcacgattt  acttcttata  aataagtgca  aatcgtcgcg  caccatatca
301 acttttggca  gccattgcac  aacaattttt  cacttagaat  atctacacct  atttttaact
361 gtagatccca  ataataact  gtccttatag  caatgggtatc  aacctacca  gaatcagagg
421 ttactctagg  aaggtaacct  tttgagcgac  tccaccaatt  gaaagtggac  accattttcg
481 gcttgccggg  tgacttcaac  ctttccttat  tggacaaagt  gtatgaagtt  ccggatatga
541 ggtgggctgg  aaatgccaac  gaattgaatg  ctgcctatgc  tgccgatggt  tactccagaa
601 taaagggatt  gtcttgcttg  gtcacaactt  ttggtgttgg  tgaattgtct  gctttaaacg
661 gagttgggtg  tgccatgct  gaacacgtag  gacttctaca  tgctgttggg  gttccatcca
721 tatcgtcaca  ggctaaacag  ttggtgctcc  accatacctt  gggtaatggt  gacttcactg
781 tttttcacag  aatgtccaat  agcatttctc  aaactacagc  atttctctca  gatattctta
841 ttgaccagg  tcaaatagat  agatgcatca  gagaagcata  tgttcatcag  agaccagttt
901 atgttggttt  accggcaaat  atggttgatc  tcaagggtcc  ttctagtctc  ttagaaactc
961 caattgattt  gaaattgaaa  caaatgatac  ctgaagctca  agaagttggt  gaaacagtcc
1021 tgaagttggt  gtcccaagct  acaaacacca  ttatcttggg  agacgcttgt  gccctcagac
1081 acaattgcaa  agaggaagtc  aaacaattgg  ttgatgccac  taattttcaa  gtctttacaa
1141 ctccaatggg  taaatctggt  atctccgaat  ctcatccaag  attgggctgt  gtctatgtcg
1201 ggacaatgtc  gagtcctcaa  gtcaaaaaag  ccgttgaaaa  tgccgatctt  atactatctg
1261 ttggttcgtt  gttatcggac  ttcaatacag  gttcattttc  atactcctac  aagacgaaga
1321 atgttggtga  attccactct  gactatatga  aaatcagaca  ggccaccttc  ccaggagtgc
1381 aaatgaaaga  agccttgcaa  cagttgataa  aaagggtctc  ttcttacatc  aatccaagct
1441 acattcctac  tcgagttcct  aaaaggaaac  agccattgaa  agctccatca  gaagctcctt
1501 tgaccaaga  atatttgtgg  tctaaagtat  ccggtctggt  tagagagggt  gatattatcg
1561 taaccgaaac  tggtagatct  gctttcggaa  ttattcaatc  ccattttccc  agcaacacta
1621 tcggtatatc  ccaagtcttg  tggggctcaa  ttggtttcac  agtaggtgca  acagttggtg
1681 ctgccatggc  agcccaggaa  atcgacccta  gcaggagagt  aattttgttc  gtcggtgatg
1741 gttcattgca  gttgacggtt  caggaaatct  ctacgttgtg  taaatgggat  tgtaacaata
1801 cttatcttta  cgtgttgaa  aatgattggt  accatataga  aaggttgatc  caccgcaaaa
1861 gtgccagcta  caacgatata  cagccttgga  accatttatc  cttgcttcgc  ttattcaatg
1921 ctaagaaata  ccaaatgtc  agagtatcga  ctgctggaga  attggactct  ttgttctctg
1981 ataagaaatt  tgcttctcca  gataggataa  gaatgattga  ggtgatgtta  tcgagattgg
2041 atgcaccagc  aaatcttggt  gctcaagcaa  agttgtctga  acgggtaaac  cttgaaaatt
2101 gaattaaaca  taccaattca  ccgctattta  gttattatat  tcatagcagc  agttatagaa
2161 ttggcttatt  atgtgttcgt  agtaattgag  tgagctgttt  catttattgt  acttatagct
2221 agaaagagaa  ccaggacgtt  actaaaacga  ctaaaactgt  tcaaagtaaa  atatattgaa
2281 aattccattc  agaatgaatt  ttacc
```

//

Revised: July 5, 2002.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Oct 3 2002 17:48:23



## Taxonomy

# Clear

## Details

## Get Subsequence

## Links

LOCUS	AF098293	2525 bp	DNA	linear	PLN 06-APR-1999
-------	----------	---------	-----	--------	-----------------

DEFINITION *Aspergillus oryzae* pyruvate decarboxylase (pdcA) gene, complete cds.

ACCESSION AF098293

VERSION AF098293.1 GI:4323052

## KEYWORDS

SOURCE            *Aspergillus oryzae*.

ORGANISM *Aspergillus oryzae*

Eukaryota; Fungi; Ascomycota; Pezizomycotina; Eurotiomycetes;  
Eurotiales; Trichocomaceae; mitosporic Trichocomaceae; Aspergillus.

REFERENCE 1 (bases 1 to 2525)

AUTHORS    Lee,D.W., Koh,J.S., Kim,J.H. and Chae,K.-S.

TITLE Cloning and nucleotide sequence of one of the most highly expressed genes, a pdcA homologue of *Aspergillus nidulans*, in *Aspergillus oryzae*

JOURNAL Biotechnol. Lett. 21, 139-142 (1999)

REFERENCE 2 (bases 1 to 2525)

AUTHORS    Lee, D.W., Koh, J.S., Kim, J.H. and Chae, K.-S.

TITLE Direct Submission

JOURNAL Submitted (13-OCT-1998) Faculty of Biological Sciences, Chonbuk  
National University, Dukjin-dong 1-ga, 664-14, Chonju, Chonbuk  
561-756, Republic of Korea

FEATURES	Location/Qualifiers
----------	---------------------

```
source      1..2525
```

```
/organism="Aspergillus oryzae"
```

```
/db_xref="taxon:5062"
```

gene <282..>2182

```
/gene="pdcA"
```

```
mRNA      join(<282..369,404..484,566..1790,1846..2053,
              2153..>2182)
```

```
/gene="pdcA"
```

```
/product="pyruvate decarboxylase"
```

```
CDS      join(282..369,404..484,566..1790,1846..2053,2153..2182)
```

```
/gene="pdcA"
```

```
/codon_start=1
```

```
/product="pyruvate decarboxylase"
```

```
/protein_id="AAD16178.1"
```

```
/db_xref="GI:4323053"
```

```
/translation="MSLSTSSGDFVRLAFVQYTVFLVSINAPAWDYNLVALDYLPKCD
LHWVGNCNELNAGYAADGYARINGMSALVTTFGVGELSALNAIAGAYSEFVPIVHIVC
QPHTKSQKDGMLLHHTLGNNGDFNVFTRMSADISCTLGCLNSTHEVATLIDNAIRECWI
RSRPVYISLPTDMVTKKIEGERLDTPLDLSLPPNDPEKEDYVVDVVLKYLHAAKKPVI
LVDACAIRHRVLDEVHEFVEKSGLPFVAPMGKGAVDETHKNYGGVYAGTGSNPGVRE
QVESSDLILSIGAIKSDFN'TTGF SYRIGQLNTIDFHSTYVRVRYSEYPDINMKGV LQK
IVQRMGNLNVGPVSPPSNLLPDNEKASTEQAITHAWLWPTV GQWLKEKD VVITETGTA
NFGIWDTRFPAGVTAISQVLWGSIGYSVGACQGAALAAKEQGRRTVLFVGDGSFQLTL
OEVSTMIRNNLNP IIFVICNEGYTIERYIHGWEAVYNDIOPWDFLNIPVAFGAKDKYK
```

GYKVTTRDELRELRFANEEFASAPCLQFEIDSRIGR"

BASE COUNT	659 a	600 c	587 g	679 t
ORIGIN				
1	aagcttgttaa	ttttggcgta	ctatctgtac	tatctacca
61	gtatggcctg	gtcttactgt	tgcacgtcga	gttgtacagg
121	tataactgcc	cccccttgtc	catgtttcca	ggaagaattc
181	aacaacctta	tcactacta	taccttccat	tttccattcc
241	ggcgacagat	atcgctacaa	gggatcttcg	caagcccata
301	caggcgactt	cgtgagggtg	gcgttcgtgc	agtacacggg
361	cgccagcttg	taccgcaact	actgacagta	tcaattcgca
421	gctttggatt	acctgccaaa	atgcgatctt	cattgggtag
481	gccggtatgt	ctccactgaa	tgggtcaatgg	gaattctggg
541	tatcgctaac	ggtgggtcaa	tgcaggatac	gctgctgatg
601	atgtctgctt	tagtcaccac	ctttgggtgtg	ggtgagctat
661	ggtgcatact	ccgaatttgt	gcctatcggt	cacattgttg
721	cagaaagatg	gaatgctcct	ccaccacacc	ttgggcaacg
781	agaatgagtg	ccgacatctc	ttgcacactt	ggatgtttga
841	accctcattg	ataatgctat	ccgagaatgt	tggattcgta
901	ctccctaccg	atatggtgac	aaagaaaatc	gagggagAAC
961	cttagtctac	caccgaacga	tcccgaaaaa	gaagattacg
1021	tatctgcacg	ctgcaaagaa	acccgttatt	cttgtcgatg
1081	gtgctcgatg	aagttcatga	gttcgtggaa	aaatctgggc
1141	atgggtaaag	gagcagtggg	tgagactcac	aagaactacg
1201	ggatcaaacc	caggtgttcg	tgagcaagtc	gaatcttcag
1261	gctatcaagt	ccgatttcaa	cacgactggg	ttctcttacc
1321	attgacttcc	atagtacata	cgtgcgcgtc	cgggtactccg
1381	aaaggcgctc	ttcaaaagat	tgttcaaaga	atgggcaatc
1441	ccgccgtcga	acctactgcc	ggacaacgag	aaggcatcaa
1501	gcatggctct	ggcctactgt	cgggcagtg	ctgaaagaaa
1561	accggcactg	ccaatttcgg	tatctgggac	actcggttcc
1621	agtcaggttc	tttggggtag	tatcggtat	tcagttggag
1681	gccgcaaaag	agcagggccg	acggactgta	cttttcgtgg
1741	acgctccagg	aagtcagcac	catgataaga	aataacctta
1801	cctttgggtg	acacatgtct	catgcccgac	taacatcgta
1861	cgaaggatat	accatcgaac	ggtacattca	tggatgggaa
1921	gccctgggac	ttcttgaaca	ttcctgtggc	attcggcgcg
1981	caaggtcaca	acccgagacg	agttgagggg	gcttttcgca
2041	accctgtctc	caggtagggt	tcatttcgtt	tacctctgtc
2101	aaccaaagca	gttggttgag	ctccacatgc	ctcgcgacga
2161	tgacagccga	atcggccgct	gagcggaaca	agtcccttta
2221	aggatagacc	agatctcttt	ccttctttac	ggctcgggaa
2281	caccatgcac	acggggcatt	tttaattcca	tcatgacatg
2341	tttgcgatag	ttttataatg	agttctttat	ctgctttccc
2401	accttctaag	ttaaaacaat	aatcagattt	aattaaataa
2461	tagagagaga	gacagagaaa	aagagaaaaa	gaaataacca
2521	tgaac			

//

Revised: July 5, 2002.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Oct 3 2002 17:48:23



PubMed

Nucleotide

Protein

Genome

Structure

PopSet

Taxonomy

Search  for 

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

    ☐ 1: AI728334. BNLGHi10479 Six-d...[gi:5047186]

Links

**IDENTIFIERS**

**dbEST Id:** 2637583  
**EST name:** BNLGHi10479  
**GenBank Acc:** AI728334  
**GenBank gi:** 5047186

**CLONE INFO**

**Clone Id:** (5')  
**DNA type:** cDNA

**PRIMERS**

**Sequencing:** T3 Primer  
**PolyA Tail:** Unknown

**SEQUENCE**

GAGAAAGCAATCATAGTACAACCAAATCGTGTGACGATAGGCAATGGTCCTTCTTTTGGC  
TGGGTTTTTCATGGCTGACTTCTTAAGTGCATTGGCAAAGAACTTAAGAAAAATACCACT  
GCCGTGGAGAACTATCGGCGAATCTTTGTCCCTCCAGGCATGCCTTTGAAGCATGGGAAT  
GACGAGCCTCTGAGAGTCAATGTTCTTTTTAAGCACATTCAGGATATGCTAAGCAAAGAC  
TCTGCGGTTATTGCTGAAACTGGAGACTCATGGTTCAATTGTCAGAAGCTCCATCTTCCA  
GAGAACTGCGGTTATGAATTCCAGATGCAGTATGGATCCATTGGTTGGTCAGTTGGTGCC  
ACGCTTGGATATGCTCAGGCAGCTAAAGGCAAACGTGTGATTGCTTGCATTGGCGATGGT  
AGTTTCCANGTGACAGCTCAAGAATATTCCAACAATGATCCCATGTGGACAAAAAAGCAT  
CATATTTCTTATCAATAATGGAGGCTATACAATTGAAGTTGAGATTCATGATGGCCCTTA  
CAATGTTATCAAGAACTGGAATTACACTGCCTTTGTTGATGCCATCCACAATGGTGAAGG  
CAAATGCTGGACAGCCAAGGTGAGGACCGANGATGAACTGATAAATGC'T

**Entry Created:** Jun 11 1999  
**Last Updated:** Jun 11 1999

**PUTATIVE ID** Assigned by submitter  
PYRUVATE DECARBOXYLASE ISOZYME 1 (PDC)

**LIBRARY**

**Lib Name:** Six-day Cotton fiber  
**Organism:** Gossypium hirsutum  
**Cultivar:** Acala Maxxa  
**Tissue type:** immature fiber  
**Develop. stage:** Six days post anthesis  
**Lab host:** XL1-Blue  
**Vector:** pBluescript II KS+

**SUBMITTER**

**Name:** Ben Burr  
**Lab:** Biology Department  
**Institution:** Brookhaven National Laboratory

Address: Upton, NY 11973, USA  
Tel: 516-344-3396  
Fax: 516-344-3407  
E-mail: [burr@bnlux1.bnl.gov](mailto:burr@bnlux1.bnl.gov)

**CITATIONS**

Title: ESTs from developing cotton fiber  
Authors: Blewitt, M., Matz, E.C., Davy, D.F., Burr, B.  
Year: 1999  
Status: Unpublished

**MAP DATA**

---

Revised: July 5, 2002.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Oct 3 2002 17:48:23



PubMed

Nucleotide

Protein

Genome

Structure

PopSet

Taxonomy


 Search  for   

Limits

Preview/Index

History

Clipboard

Details

     
☐ 1: Z54096. *S.pombe* chromosom...[gi:984221]

Links

LOCUS SPAC13A11 12168 bp DNA linear PLN 18-OCT-1999  
 DEFINITION *S.pombe* chromosome I cosmid c13A11.  
 ACCESSION Z54096  
 VERSION Z54096.1 GI:984221  
 KEYWORDS cyp51; cytochrome P450; cytosol aminopeptidase; Fes/CIP4 homology domain; pyruvate decarboxylase; Rho GTPase protein; Thiamine pyrophosphate enzymes; ubiquitin carboxy-terminal hydrolase.  
 SOURCE fission yeast.  
 ORGANISM *Schizosaccharomyces pombe*  
 Eukaryota; Fungi; Ascomycota; Schizosaccharomycetes; Schizosaccharomycetales; Schizosaccharomycetaceae; Schizosaccharomyces.  
 REFERENCE 1 (bases 1 to 12168)  
 AUTHORS Hunt,S., Devlin,K., Churcher,C.M., Barrell,B.G., Rajandream,M.A. and Walsh,S.V.  
 TITLE Direct Submission  
 JOURNAL Submitted (06-SEP-1995) *Schizosaccharomyces pombe* chromosome I sequencing project, Sanger Centre, Hinxton Hall, Hinxton, Cambridge CB10 1RQ E-mail: barrell@sanger.ac.uk  
 COMMENT Notes:  
 Details of yeast sequencing at the Sanger Centre are available on the World Wide Web.  
 (URL, [http://www.sanger.ac.uk/Projects/S\\_pombe](http://www.sanger.ac.uk/Projects/S_pombe))  
 Protein coding regions (CDS) have been predicted with the help of computer analysis using the Genefinder program in PomBase (an ACEDB database) with additional predictions for the branch-acceptor sites supplied by the program Sp3splice. CAUTION: It is possible that for any individual CDS we may have underestimated or overestimated the number of introns/exons or we may not have chosen the correct splice donor/acceptor sites.  
 CDS are numbered using the following system eg SPAC5H10.01c. SP (*S. pombe*), A (chromosome 1), c5H10 (cosmid name), .01 (first CDS), c (complementary strand).  
 The more significant matches with motifs in the PROSITE database are also included but some of these may be fortuitous.  
 The length in codons is given for each CDS.  
 IMPORTANT: This sequence MAY NOT be the entire insert of the sequenced clone. It may be shorter because we only sequence overlapping sections once, or longer, because we arrange for a small overlap between neighbouring submissions.  
 Cosmid c13A11 overlapped at the 5' end by cosmid c2F7 and at the 3' end c3H8.  
 FEATURES  
 source Location/Qualifiers  
 1..12168  
 /organism="Schizosaccharomyces pombe"  
 /strain="972h-"  
 /db\_xref="taxon:4896"



/chromosome="I"  
/map="IL"  
/clone="cosmid c13A11"  
gene complement(join(1..340,385..487))  
/gene="SPAC13A11.01c"  
/note="SPAC2F7.18c"  
CDS. complement(join(1..340,385..487))  
/partial  
/gene="SPAC13A11.01c"  
/note="SPAC13A11.01c, len:146, SIMILARITY:Saccharomyces cerevisiae, YFE7\_YEAST, hypothetical 82.2 kd protein in EMP47-SEC53 intergenic region, (714 aa), fasta scores: opt: 263 E():2.3e-09, (29.5% identity in 146 aa overlap)"  
/codon\_start=1  
/product="putative Rho GTPase protein"  
/protein\_id="CAA90802.1"  
/db\_xref="GI:984222"  
/db\_xref="SWISS-PROT:Q09697"  
/translation="MISSFSNGFWSKDYATGVKKLFDCLDNGVEENEQVKNLLKLYKE ANEEFGEKLQEITKECLKGKKPENTEDGATSNKAFEGLRSEIANQGKQHIRIAKDLET LIIAPFSKMSIDHSQKLQTSQQVLTNQIKSYEKYYTLKKTSAAY"  
misc feature 1..104  
/note="overlap with cosmid c2F7, positions 42407 42510 EM:Z50142 S.pombe chromosome I cosmid c2F7"  
misc feature complement(join(144..340,385..469))  
/gene="SPAC13A11.01c"  
/note="Match to PF00611 FCH, Fes/CIP4 homology domain Score 103.18"  
misc feature complement(341..354)  
/gene="SPAC13A11.01c"  
/note="ctaacaacgaatag, splice branch and acceptor"  
misc feature complement(379..384)  
/gene="SPAC13A11.01c"  
/note="gtattt, splice donor sequence"  
gene complement(1251..2738)  
/gene="SPAC13A11.02c"  
/note="cyp51"  
CDS complement(1251..2738)  
/gene="SPAC13A11.02c"  
/note="SPAC13A11.02c, len:495"  
/codon\_start=1  
/product="probable cytochrome p450 51"  
/protein\_id="CAA90803.1"  
/db\_xref="GI:984223"  
/db\_xref="SWISS-PROT:Q09736"  
/translation="MAFSLVSILLSIALAWYVGYIINQLTSRNSKRPPIVFHWIPFVG SAVAYGMDPYVFFRECRAKYGDVFTFVCMGRKMTAFLGVQGNDFLFNGKLADLNAAEEA YSHLTTPVFGKDVVDIPNHVFMEHKKFIKSGLGFSQFRSYVPLILNEMDAFLSTSPD FPGKEGVADLLKTMPTMTIYTASRTLQGAEVKRGFDAGFADLYHDLDQGFSPVNFVF PWLPLPRNRRRDRAHKIMQKTYLKI IKDRRSSTENPGTDMIWTLMSCKYRDGRPLKEH EIAGMMIALLMAGQHTSAATIVVWLALLGSKPEIIEMLWEEQKRVVGENLELKFDQYK DMPLLNVIQETLRLHPPIHSHMRKVKRDLVPVPGSKIVI PANNYLLAAPGLTATEEEY FTHATDFDPKRWNDRVNEDENAEQIDYGYGLVTKGAASPYLPFGAGRHRICIGEQFAYM HLSTIISKFVHDYTWTLIGKVPNVDSMVALPLGPVKIAWKRRN"  
misc feature complement(1263..2642)  
/gene="SPAC13A11.02c"  
/note="Match to PF00067 p450, Cytochrome P450 Score 169.72"  
misc feature complement(1407..1436)  
/gene="SPAC13A11.02c"

/note="PS00086 Cytochrome P450 cysteine heme-iron ligand signature"  
gene join(4637..4702,4797..5363)  
/gene="SPAC13A11.03"  
CDS join(4637..4702,4797..5363)  
/gene="SPAC13A11.03"  
/note="SPAC13A11.03, len:210, SIMILARITY:Arabidopsis thaliana, CAB43931, hypothetical 14.7 kd protein., (128 aa), fasta scores: opt: 236, E():2.7e-08, (37.5% identity in 120 aa)"  
/codon\_start=1  
/label=SPAC13A11.03  
/product="hypothetical coiled-coil protein"  
/protein\_id="CAA90804.1"  
/db\_xref="GI:984224"  
/db\_xref="SWISS-PROT:Q09739"  
/translation="MPPKGLSLAEKRRRLEAIFHDSKDFQKKEVEKLGSKKQIVLQTVKDVLQSLVDDNIVKTEKIGTSNYYWSPSDAKRSRESVLGSLQAQLDDLKQKSKTLDENISFEKSKRDNEGTENDANQYTLELLHAKESLKLKTKQLSNLHNCPETFELKNENTKKYMEANLWTDQIHTLIAFCRDMGADTNQIREYCSIPEDLDDLQLPIL"  
misc\_feature 4703..4708  
/gene="SPAC13A11.03"  
/note="gtaagt, splice donor sequence"  
misc\_feature 4775..4796  
/gene="SPAC13A11.03"  
/note="ctaactctatgttttttttatag, splice branch and acceptor"  
gene complement(join(5836..5905,5951..6015,6069..6366,6412..7099,7200..7332,7411..7506))  
/gene="SPAC13A11.04c"  
CDS complement(join(5836..5905,5951..6015,6069..6366,6412..7099,7200..7332,7411..7506))  
/gene="SPAC13A11.04c"  
/note="SPAC13A11.04c, len:449, SIMILARITY:Schizosaccharomyces pombe, O74442, ubiquitin carboxyl-terminal hydrolase, (457 aa), fasta scores: opt: 536, E():3.1e-27, (30.6% identity in 320 aa)"  
/codon\_start=1  
/label=SPAC13A11.04c  
/product="ubiquitin carboxyl-terminal hydrolase"  
/protein\_id="CAA90805.1"  
/db\_xref="GI:984225"  
/db\_xref="SWISS-PROT:Q09738"  
/translation="MPGDVEGCQHLKLKPADVENYQKICTQIFSCHFVPRRCSTCKRI NKRSIRCLSCHSVGCLWGHGEEHAMEHTHMIGVDVKNGHTYCFGCQDYVYQTELETL RFKIKNIKAWQSDHKRLPEKYNQMVCLEAYRKYPVCATAGLRGIQNLGATCFMSVIL QSILHNPLVRNLFFSGFHTSTDCKRPTCMTCAIDDMFSSIYNSKNKSTFYGPVAVLNL MWKLSKSLCGYSQQDGHEFFVYLLDQMHTESGGGTSMPCTCPIHRIFSGSLKNVVTCL DCKKERVAVDPLMDISLDINEPTLQGCLERFVSKEKVQYSCHSCGSKNAIKQLVFDKL PPTICMQLKRFEQNNFAMSTKIDKQVSYPAFLRMRYNFNQDDVDYQLYSVVCHKGTLD TGHYIAYTYQYQNWFLDDTTIVEVKESEVLNSQAYLLFYHERQILYSDEMTVKTEN"  
misc\_feature complement(5906..5920)  
/gene="SPAC13A11.04c"  
/note="ctaaccagattatag, splice branch and acceptor"  
misc\_feature complement(5945..5950)  
/gene="SPAC13A11.04c"  
/note="gtaggc, splice donor sequence"  
misc\_feature complement(6016..6028)  
/gene="SPAC13A11.04c"  
/note="ctaattttttag, splice branch and acceptor"  
misc\_feature complement(6063..6068)

```

/ gene="SPAC13A11.04c"
/ note="gtatat, splice donor sequence"
misc_feature complement(6069..6149)
/ gene="SPAC13A11.04c"
/ note="Match to PF00443 UCH-2, Ubiquitin carboxyl-terminal
misc_feature hydrolases family 2 Score 53.18"
complement(6367..6385)
/ gene="SPAC13A11.04c"
/ note="ttaacaaatattatttttag, splice branch and acceptor"
misc_feature complement(6406..6411)
/ gene="SPAC13A11.04c"
/ note="gtaaaa, splice donor sequence"
misc_feature complement(7018..7071)
/ gene="SPAC13A11.04c"
/ note="Pfam match to entry PF00442 UCH-1, Ubiquitin
misc_feature carboxyl-terminal hydrolases family 2"
complement(7100..7110)
/ gene="SPAC13A11.04c"
/ note="ctaattcttag, splice branch and acceptor"
misc_feature complement(7194..7199)
/ gene="SPAC13A11.04c"
/ note="gtacgt, splice donor sequence"
misc_feature complement(7261..7278)
/ gene="SPAC13A11.04c"
/ note="PS00190 Cytochrome c family heme-binding site
misc_feature signat ure"
complement(7333..7352)
/ gene="SPAC13A11.04c"
/ note="ttaacatttctttcatttag, splice branch and acceptor"
misc_feature complement(7405..7410)
/ gene="SPAC13A11.04c"
/ note="gtaagg, splice donor sequence"
gene 8608..10149
/ gene="SPAC13A11.05"
CDS 8608..10149
/ gene="SPAC13A11.05"
/ note="SPAC13A11.05, len:513, SIMILARITY:Bos taurus,
AMPL_BOVIN, cytosol aminopeptidase, (487 aa), fasta
scores: opt: 1008, E():0, (42.1% identity in 442 aa)"
/ codon_start=1
/ label=SPAC13A11.05
/ product="cytosol aminopeptidase"
/ protein_id="CAA90806.1"
/ db_xref="GI:984226"
/ db_xref="SWISS-PROT:Q09735"
/ translation="MKGLGLSTRTFNWSSLSSILLPRIPLATTKADSLILAVRHDKQV
FSEDYRQVVDQYFETSPKKNDIRLFWNTQGFVRLAIVQLEENVSEKSVRSAAAEAAKI
LKSNAGKSIADVGMGFPKDAALGAALATYDFSLRRDHLSVYQDEKVVKEKNLFTSPAP
ERLTFQLLSNTSEKKTATAEENAFKVGLIEAAQNLARSLMECPANYMTSLQFCHFAQ
ELFQNSSKVKVFVHDEKWIDEQKMNGLLTVNAGSDIPPRFLEVQYIGKEKSKDDGWLG
LVGKGVTFDSGGISIKPSQNMKEMRADMGGAAVMLSSIYALEQLSIPVNAVFTPLTE
NLPSGSAAPGDVIFMRNGLSVEIDNTDAEGRILADAVHYVSSQYKTKAVIEASTLT
GAMLVALGNVFTGAFVQGEELWKNLETASHDAGDLFWRMPFHEAYLKQLTSSSNADLC
NVSRAAGGCCTAAAFIKCFLAQKDLFAHLDIAGVMDKQLNSWDCDGMGRPVRTIIE
VARKY"
misc_feature 8743..10047
/ gene="SPAC13A11.05"
/ note="Match to PF00883 Peptidase_M17, Cytosol
aminopeptidase family Score 764.41"
misc_feature 9685..9708

```

/gene="SPAC13A11.05"  
 /note="PS00631 Cytosol aminopeptidase signature"  
 11713..12168  
 /gene="SPAC13A11.06"  
 /note="SPAC3H8.01"  
 11713..12168  
 /partial  
 /gene="SPAC13A11.06"  
 /note="SPAC13A11.06, len:151,  
 SIMILARITY:Schizosaccharomyces pombe, O42873, putative  
 pyruvate decarboxylase, (570 aa), fasta scores: opt: 709,  
 E():0, (71.2% identity in 153 aa overlap)"  
 /codon\_start=1  
 /product="pyruvate decarboxylase"  
 /protein\_id="CAA90807.1"  
 /db\_xref="GI:984227"  
 /db\_xref="SWISS-PROT:Q09737"  
 /translation="MSGDILVGEYLFKRLEQLGVKSILGVPGDFNLALLDLIEKVGDE  
 KFRWVGNTNELNGAYAADGYARVNGLSAIVTTFGVGELSAINGVAGSYAEHVPVWHIV  
 GMPSTKVQDTGALLHHTLGDGDFRTFMDMFKKVSAYSIMIDNGNDAAEKI"  
 misc\_feature 11755..12162  
 /gene="SPAC13A11.06"  
 /note="Match to PF00205 TPP\_enzymes, Thiamine  
 pyrophosphate enzymes Score 168.62"  
 misc\_feature 12046..12168  
 /gene="SPAC13A11.06"  
 /note="615 (+1) 1 123 SPAC3H8 EM:Z69086 S. pombe  
 chromosome 1"

BASE COUNT 3926 a 2206 c 2256 g 3780 t

ORIGIN

```

1 tagtaagcgc tttttgtttt ttttaagtgt tagtatttct tttcataact ctttatttga
61 ttggtaagaa cttgttggct ggtttgtaac ttctgagaat gatcaataga catttttagag
121 aaaggagcaa taattagcgt ttctaagtc ttggcaatac gaatatgttg ctttccttga
181 ttagcaatct cggagcgcaa cccttcaaaa gccttgtag aagtagcacc gtcttctgta
241 ttctctggct ttttcccttt tagacattct ttcgttattt cttgaagctt ttcaccaa
301 tcctcattcg cttctttata taacttcaac aagtttttta ctattcggtt ttagaggaag
361 tcctatatta atttatttaa ataccttgct cattctcttc aactccatta tccaggcaat
421 cgaaaagttt tttcacacca gtgcgtagt ctttactcca aaaaccattt gaaaagctcg
481 atatcatttc aattcgagac tgaaactgtc acaacctgtc aagaaatata gaggtggttg
541 acatgtctac gccctttatt actcaactta agtaaatagc gaattgctac gcagaaaagt
601 tttaccatc aaggatgcca aaagtatcca gtaaataaga caacagatac gttactgac
661 acacataaga tagcatacca ctaactcctt ataaaatgta aacttcattc agttgaatag
721 cagcctacgc ttttaccac tttgtattat taatgctatg gttggatcga tgggtacaatt
781 aaaaactaca aaattttaat taatagctta ctgagaaacc ttgcgtttgc acgatgaggt
841 tttgaaaaga gagagtagcg tactttattg ttataatgcc tacatgaaaa cgaaagagag
901 cgtcccgacg ttattaaaac tataatatcg taaaaatgta tactaatga tgacagggta
961 gttttcatat cgatttagtg cgaggataag gaagtttgaa attgcataat acattctttt
1021 gacatttgaa atgctaccaa atagtcattt aaaacgctaa aatatttaca atttcataaa
1081 acaataaagc acttaaaatt taaaatgtct agcaaatttc atggttattc gttactcgtc
1141 agataaaaca aaaacgttta acacgggcat gaaaagagaa aggcggaacc aaagttagaa
1201 accaatacca taagcgtatt gcaaatgcat tcaaattagg ggaaattccc ttaattacga
1261 cgcttccaag caattttaac aggacccaag ggaagagcaa ccatactgga atagtcaacg
1321 ttgggaactt ttccaataag agtccaggtg taatcgtgta caaacttgct aataatagtg
1381 ctcaaagca tgtaggcaaa ttgctcacca atgcaacgat gacgaccagc tccaaagggt
1441 aggtatggag aagcagcacc ctttgtaacg agaccataac cataatcaat ttgctcagca
1501 ttttcgtctt cattgacgcg gtcattccag cgcttggggg caaagtcagt agcatgagtg
1561 aagtattcct cttcagtagc agttaaacg ggagcagcaa gaagatagtt gttggcaggg
1621 ataacaatct ttgaaccagg gacgggtaag tcacgtttga ctttacgcat atgactgtgg
1681 atagggggat ggagtcctaa agtttcttga ataacatagt tgagaagagg catgtcctta
1741 tattggtcaa acttcaattc caagttttca cctacaacac gcttctgctc tccccacaac

```

```
1801 atttcaatga tttcaggttt gctacccaaa agagcaagaa cccaaacgat agtggcagca
1861 gaggtgtgtt gaccagccat caaaagtga atcatcatat cagcaatttc atgttccttc
1921 aaaggacggc catctctata cttagcagtc atcaaagtcc aaatcatatc agtaccagga
1981 ttttcagtgc tagagcgacg atctttaata attttaagat aagttttttg cataattttg
2041 tgggctctat cacggcgtct gtttctaggg agaggtaacc aagggaatac aaagttaaca
2101 gggctaaatc cttgatcaag gtcattgtac aagtcagcga aaccagcgtc aaagcccttg
2161 cgaacctcag caccttgtaa agtacgagaa gcagtgtaaa tagtcatgac aggcattggtc
2221 ttcagcaaat cagcaacgcc ttccttacca ggtccgaagt caggagaagt gctgagaaaa
2281 gcatccattt cgttaagaat gagggggaca taagagcggg attgagagaa gccaagacca
2341 gatttgataa acttcttgtg ctccataaaa acatggttgg gaatatcgta aacgacatct
2401 ttaccgaata caggggtagt taaatgagag taagcttctt cagcattcaa atcagctaata
2461 ttaccgttaa agaggaaatc attaccttgg acaccaagga aagcgggtcat tttacgaccc
2521 atgcaacaaa aggtgaaaac atcaccatat ttggcagcag actcgcgaaa aaatacataa
2581 gggctccatac cgtatgcgac agcagaacct acaaagggta tccaatggaa gacaattggg
2641 gggcgtttgc tattgcgaga ggtaagtggg ttaataatat aaccaacata ccaagccaat
2701 gcaatcgata ataaaatcga aactaaagag aatgccatcc tgacttatga aaaaaaaagt
2761 aatgctcaag tgggtaaagc aggtaaaaaa taatcagacg aaaaaataa aatcagtcag
2821 taaagtagta aggggtaagc agatagcaac caaaaacgaa taagaccaac caaaaaacaa
2881 cgacaacaac aacaaatgtg taacagataa aaatagttca acccttttag ctaatgcaaa
2941 taaacttgaa aaagaaaagg agaaagaaaa aatgaattga ttaacacgca gctacgagaa
3001 caacgaatga aacaaataga aaaaagcgtt aaaaacgaaa tcgaaaacag acaataatat
3061 gaaagaaaaa caaaaaaaa ttacgtttcc tttcacttaa cgaatgagat tgggattttg
3121 aaaagacaat tttcaaaaac aacgatcgat acagaaattt tagatcttgt cttttttaag
3181 gtacttctac aagtgaagaa tcgatttata ttctcttga caccataatc gctggtttcg
3241 acgacgattc aagctaactt gggctctcgc ctgactgatg ggagttggta gaattatata
3301 tgcaacataa tggcgtgtga accgtttggt tatgctgaaa atccgtaaaa aaacatgcta
3361 cttgatcact tcaaaattac acttttttat ccaatcatat ttttcagata ctaccttatt
3421 gttctttctg gtcaaggtga attagccact aagccaatga gaaatccaat tttgttacct
3481 gagaataggg cgacatcgta cgattcgagg gtttgtgagg tgaatgactg cggggcttat
3541 gttgcgatcg aacgatggca cgtccattgg accgagataa tacacatgtt taccgtagct
3601 cgtaacgcac atttaacaaa ctacttgctt ctaccctact ttatcaccta aaaataaaaa
3661 tgtgtcaaca tgtgcacgtc tactatacct tctactattt gttcggtaaa ccaccaaccg
3721 tgccctcctta tacgtcttct ccaaacatgt aattggattc gaacaacttt agctcgacaa
3781 gtcgttttagc ttacgaacat ccaaagctt atctaatac ctaactaact cgctcttttt
3841 ggctcgcatga ctttgcatatc gattgtttag taataaaaaa aataatcctt ttcacacctg
3901 tttttaccca tgggaaagcg gttcccaaag tttgtcactt actcccatat tctaattctag
3961 attaaaacac atttagctgc gaacatggca caccgggtta taccagacca tagttgaaat
4021 tagtagacaa attgactgag tatctcttat caaaagtcaa tacacatttt gtttgcgctg
4081 acattaaaca tcgttgctct tatgaatgct atgtgtgtgt gtgtgtgtat gtggaaatgt
4141 gttagtcaag cagtgtttgc ttgaaattcg ctgtgtgaaa ataataaagt gaaaaataaa
4201 taaaagggtt taaaaccgaa ttctgtacag aagggtgagc cgcacaaaag ctcttgagaa
4261 aaaaaaatc atgctgctat cagtattcga gtcttttcga tttcgtggca aaaggaagtt
4321 ttctctacat gtttttgtgc gacaatagcc aacccattc acttaattcc ttaatttctt
4381 caagggtaaa aaagattagg atagctgaat aacgttttgc attgcattat gttgcaatac
4441 gatagtaaga cattataggg tatatatcat agttggaaag tgcggatgtg catttttagca
4501 gactacgtta tcattcagta aatgtttggg atctaaactc tcatttctcc aaaagattag
4561 aaagcatcta cttcttgcac cacaacaaac ttttaaactc catatttggt caaactaatt
4621 aataaatttt taaacaatgc ctcccaaggg actatcgctt gcagagaaaa gaagaagatt
4681 agaggccatt ttccatgact ctgtaagttc gtgatatgat tgttcaagtg ccaaaacatg
4741 atttttgccc attcgtgtct cactttccac attgctaatac tatgtttttt ttatagaaag
4801 atttttttca actaaaggag gtggaaaagc tgggctctaa aaagcaaatac gttttgcaaa
4861 ctgtaaagga cgttttacag tctctcgttg acgacaatat cgtcaaaacg gaaaagattg
4921 gcactagcaa ttactactgg tcttttccct ccgatgctaa acgttcaaga gagtcagttc
4981 tcggatctct tcaagctcaa cttgatgatt tgaaacaaaa atctaaaact ttagatgaaa
5041 acattagttt cgagaaatcc aaaagggata acgaaggaac cgaaaatgat gccaatcaat
5101 acactttgga actgctacat gcaaaagaat cagaactaaa acttcttaaa acacaacttt
5161 caaatttaaa tcaactgtaat cctgaaacct ttgaattaaa aaatgagaat actaaaaagt
5221 acatggaagc tgccaatttg tggaccgatc aaatccatac tttgatagct ttttgcgag
5281 acatgggtgc cgataccaat caaattcgtg agtattgcag cattccagag gatttggacg
5341 atctgcaact accgattttg tgaacttgac atgcttctt aattcaaaga gttgttttcc
```

5401 ttattaatat tttaaaat ttttataacatttt aacagttaca tataaaatag cttactcgac  
5461 gacaagactt agctttttaa agataaaatt ataattagtt ggaataatat taaacgaaac  
5521 aaaacaaaag tccaaattcc tcaaaacttat agtatttggt aactttgaaa gaaaaaaagc  
5581 atgtttcttt ttttatagtt ttaacttttt ttataaattt acgtaagggtc attaatataa  
5641 atgcgactgt tgtaacggtt ccaaatccaa tgccatttct ggctcaaca atatgaaatc  
5701 taaaagcttc ttcattgagt agattatcga agtgatgatg ttccataacc acataaataa  
5761 ggaacctttg agtcaattac tttacatggt tcataatgat tgttcatgaa cggatatctg  
5821 atcgcataat ctaactcaat tttccgtttt aacagtcatt tcatcactgt ataggatttg  
5881 acgctcatgg taaaataata ggtaactata atctgggttag tatgtaaaaa aataacagca  
5941 aagagcctac gcttgagaat ttaagacctc cgatttcctt acttccacaa ttgtggtgct  
6001 atccagcaaa aaccactaca aaaattagta aaaatttcca taaaaaatc aaaaaaaatt  
6061 taatatactt gatttttggt gtatgtatag gcaatatagt gccctgtgtc gagagttcct  
6121 ttatgacaca caactgagta taattgataa tccacatcat cttgattaaa gttgtagcgc  
6181 attcttaaaa atgctggata agatacttgc ttatctattt tagttgacat ggcaaagtta  
6241 ttttgttcaa atcgtttcaa ttgcatgcaa atagttggag gtagttgtc aaagaccaat  
6301 tgtttaatat cattttttga cccgcaacta tgacagctat attgaacttt ctctttggaa  
6361 acaaacctaa aataatattt gttaaaaaat ctaaaaacga gcaaatttta cttttccaaa  
6421 catccttgaa gtgttggtc attgatgtct aaactaatgt ccatcaaagg atccacagcc  
6481 acccgctcct tcttgcaatc taggcattgta acaacgtttt tcagagatcc actaaaaata  
6541 cgatgaatat gacaagtgca cggcattgat gttcctcctc cgctttccgt atgcatttgg  
6601 tccagcaagt atacaaaaaa ttcgtggcca tcttgctggg agtaaccaca taaagatttg  
6661 ctgagtttcc acataagatt taagacagca gtgggaccgt aaaaagtaga tttgttttta  
6721 gaattataaa tagatgaaaa catatcgtca atggcacaag tcatgcaagt aggtcgctta  
6781 caatcagtag aagtatgaaa cccactgaaa aacaagttt gaaccaatgg attatgcaaa  
6841 atactttgta aaatgacgct catgaaacaa gtagcaccca agttctggat cccctcaag  
6901 cctgctgttg cgcaactgg agggattttt cgataggctt ctaagcaaac catttgattg  
6961 tacttctcgg gaagtctctt gtgataactt tgccatgcct tgataattct gatcttaaaa  
7021 cgaagggttt ccaactcggg ctgataaacg tagtcttgac agccaaagca gtaggtatgg  
7081 ccatttttta catccacgcc taagaattag taattggtat tcgggaattg catcctaact  
7141 cgatattcgc aaaaagggtt cgttgctaaa tcaatagcta tgttttcaac taaacgtacc  
7201 gatcatatga gtgtgctcca ttgctgttc ttcacatga tgtcccaaa gacatccac  
7261 agaagacaa ctcaagcatc gaatactccg cttgttgatg cgtttgcaag tactgcagcg  
7321 acgtgggaca aactaaatga aagaaatgtt aaaaggaatc gaatggaaaa agaaaattga  
7381 actgtatgaa gtgttggtca caagccttac atgacaactg aatatctggg tacagatctt  
7441 ttgatagttt tcaacatccg ccggtttaag ttaagatgc tggcaacctt ctacatcccc  
7501 aggcacctt gatagctaca aaattggcct ttcagactaa aaagcatctg cttcgatgca  
7561 gggtgaaacg acaacaatag acgaactcgc tattctggca aaagaacatt atttctctt  
7621 gcttaaat tttggcaattaa ggatgcggtg tacagcggtt agagccataa agtactcata  
7681 gtgtctgcac tactagatac cgatttcgta taaagagcta tcatagctac gctaagtcaa  
7741 ttgataaaac accaaaatag atgttcttcg tagttcttga gtgtccttg ttttctctt  
7801 taaaacataa acttggtaaa ttcccataga tttggctgtt gagtactgc aaaataaaaa  
7861 aaattactgc tagtctttaa atcacggtt aatagatgaa attgttaatt ttttaaaagc  
7921 ttggtcaaat tagactaatg cctctattaa cgtcgaacgt gtttaagtgt acgcttatgt  
7981 ttgttgccat gaaacgtttg cagggtctaaa tataaaacca tatatgacta tttaccgtga  
8041 gcatctttgt aatacaaccg agtactgca agcagggtaca tgtacgaaaa tttttgtcta  
8101 ttacatttgt atagaattta ttacttaaca attgactagg ggcagatttt tgtgccgaaa  
8161 cagtttgtgt taccctaaat agagattgta gggcaacaat acgtctgcag aagtcaaat  
8221 atatatgtag caaatgagtg tgacttactt aaaaattggg ttttgatcaa gaaataaaat  
8281 cctttggtgc tttttctttt ttatgtctag cttgttgtaa tgtattttt catcacaatg  
8341 ttgatcttgt ctactcaata acacactaga ctatgttaag tacacttgcc cttgccaat  
8401 ctgcaatcgg acatcgtatt ttatttttgc atactgaatt tacgtgtatt cgtagtttca  
8461 cgagttaacg attttogaat atccaactct cattgaactt ataataatcc gtaatcctt  
8521 acctttttt tgattacaag tgtctaatta agttttctaa atcaagagaa atcaaagata  
8581 caagcaaagc ttttttctt acgtacaatg aagggtttag gtctaagtac gagaacctt  
8641 aattggagct cactttctt cattctttta ccaagaattc ctttagctac taaaaagcc  
8701 gattccttga ttttggctgt tcgtcacgat aaacaagttt tttctgaaga ttatcgacaa  
8761 gttgttgacc agtattttga aacatctccc aaaaaaatg acattcgtct cttttggaac  
8821 actcagggtt ttgttcgtct tgccatagtc cagcttgaag aaaacgtttc tgaaaagtcc  
8881 gttcgttcag ctgctgcaga ggcagctaa attttaagt cgaatggtgc aaagtgcata  
8941 gctgtcgatg gtatgggtt tccaaggat gctgccttgg gtgtgcctt ggccacgtac

```
9001 gattttttctc ttcgtagaga ccatttgagc gtctatcaag atgaaaagggt ggttgaaaag
9061 gaaaatttat ttacctctcc tgctcctgaa aggttaactt tccaattgct tagcaacaca
9121 tctgagaaga aaactgcaac cgcagaggag aatgctttta aggtagggtt gattgaagca
9181 gctgctcaaa acttggtctcg ttctcttatg gaatgtcctg ctaattacat gacttctctt
9241 caattttgtc attttgctca agagttattc caaaattcct caaagggtta agtattcggt
9301 cacgatgaga agtggattga tgagcaaaag atgaacgggt tacttaccgt gaatgctggg
9361 tccgatattc cacctcgttt cttagaagta caatacattg gtaaagaaaa atcaaaagat
9421 gatggatggc ttggtttagt tggaaaagggt gtaacggttg acagcgggtg tatcagtata
9481 aagccttctc aaaacatgaa ggagatgcgt gctgatatgg gtggtgctgc tgttatgctt
9541 tctctatatt atgccttgga acaactttcc atccccgtga atgccgtggt tgttaccctt
9601 ttaactgaga atcttccttc tggctctgct gctaagccag gagatggtat ttttatgcgc
9661 aacggtctta gtgtagaaat tgataacact gacgctgagg gtcgtcttat tcttgccgat
9721 gcagttcact atgtctcttc tcagtataag accaaggctg tgattgaagc cagcacatta
9781 actggtgcta tgctagtagc cctcggtaat gtatttactg gtgcatttgt tcaaggagaa
9841 gagttgtgga agaactctga aactgcttcc catgatgctg gagacctttt ctggcgtagt
9901 ccatttcacg aggcttacct caagcaactt acaagttctt caaatgctga tttgtgcaat
9961 gttagtcgtg ccggtgggtg ttgttgtaga gcggctgcct ttattaaatg cttcttagcc
10021 cagaaagatt tgtcttttgc tcathtagac attgccggag ttatggataa gcagcttaat
10081 tcttgggatt gtgacggcat gagcggacgc ccagttcgca ctattattga agtcgctcgc
10141 aagtatttagc ttatttattt tcttggctat atgtgcaaac tttatgtttt tacttattta
10201 ctgttgatta caccaatctc atttttcaat gtccttggtt agtgtatgtc aatgatatgt
10261 agctatcatg attttattct ttttaatact gttttccaca accggttcata acggttgtaa
10321 ggaccgagtc tttctgatgg ctctggttct gtttcttta aaaataaaat caattcgggt
10381 gaagtctcat gttgttaa atctcttcg aagtagcgaa tttttttt ctattttgaa
10441 tatagatttg tttattaaat gtagtaatca atttactcat gactattact tttgtcaaac
10501 aaaatcataa tacaaaacta aatcttttaa ggtagaatat tctcctgtgc gaggttacaa
10561 tactttggga tatgatcaaa tcaatttttt ttagatatct gttgaaattt ttataacagt
10621 aattttgtat tatttatcat actatttatt agcgttacct gccctcttaa atgtacacat
10681 catgtggggg tgggggaacc agtgctggag atcacttctt gctataccgt ttggctcatg
10741 gcaactattg aggatagcaa ggagcgttac aatttgcaaa tcaatatgat aaacagaaaa
10801 acaacaagtt gaagcttaac actgtagcaa tataacaatg aaaaaataa tcagaattaa
10861 agcaagaagc ctataaatgg tccatcaatt gtgaaatcga tattgctaga tgtcaggcaa
10921 atcgtgaact cgggtgaaaa atgtcggttaa tattcatgtg aaatcataat tacagcaata
10981 tgttcaaaga cttttggata ctttataata atgaggattc ttttttgag tgggtcacatg
11041 gtctcgtaac catgaggtag gccctacttt tacggacgaa acaatatttc acattcagct
11101 ctcaatgtta agtatccaaa caaacatggt acaaaccaat aaagcatgta aattaataag
11161 cttgaaataa ggaatagtc ttaaattcag ttttagacata atgtaaaagg tatcaatttt
11221 gtctttggta tagatacaat taagaatttc gagacggaac gttcgggtgat gtagagtttt
11281 cataataatt tctgcagttg cttacactaa ccgaagaatc gtccagtata aactgctgac
11341 gatgccttaa atcaataatt gcaaatgca tagttgacgt tttcccggtt ggcacatagt
11401 gagatttgag tcgaataatg aaatagcctc tgacgagtgg tttcgtacca agagtttcaa
11461 aaagaataat ccacattaac tcaatgtatc attgatgacg gaatgttgc tatatagcca
11521 gagtaaccct gctctgaagc atctaagggt atatcttttt aacttttagc tagcaaaactc
11581 acttgttctt ttaagttcag aaagtatctg ctggttattc aacagataat tatcttgcca
11641 aaaaggaata cacactcagc attgtaattt gttgagtata tttcacaagc acattaaatt
11701 gcattatcaa acatgagtgg ggaattttta gtcgggtgaat atctattcaa aaggcttgaa
11761 caattagggg tcaagtcocat tcttgggtgt ccaggagatt tcaatttagc tctacttgac
11821 ttaattgaga aagttggaga tgagaaattt cgttgggttg gcaataccaa tgagttgaat
11881 ggtgcttatg ccgctgatgg ttatgctcgt gttaatggtc tttcagccat tgttacaacg
11941 ttcggcggtg gagagctttc cgctattaat ggagtggcag gttcttatgc ggagcatgtc
12001 ccagtagttc atattgttgg aatgccttcc acaaagggtg aagatactgg agctttgctt
12061 catcatactt taggagatgg agactttcgc actttcatgg atatgtttta gaaagtttct
12121 gcctacagta taatgatcga taacggaaac gatgcagctg aaaagatc
```

//

Revised: July 5, 2002.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Oct 3 2002 17:48:23



09/754,083

## Characterization of *Bacillus stearothermophilus* Plasmid pAB124 and Construction of Deletion Variants

By ALISTAIR H. A. BINGHAM,\* CHRIS J. BRUTON†  
AND TONY ATKINSON

Public Health Laboratory Service, Centre for Applied Microbiology and Research,  
Porton, Salisbury, Wiltshire SP4 0JG

(Received 18 October 1979; revised 26 January 1980)

A restriction endonuclease cleavage map of the tetracycline resistance plasmid pAB124, originally isolated from *Bacillus stearothermophilus*, was constructed using ten enzymes. Tetracycline resistance was associated with a 1.95 megadalton (Md) region of pAB124 lying between two *EcoRI* sites, and this region was circularized to produce a viable tetracycline resistance plasmid (pAB224), with two *EcoRI* fragments of pAB124 deleted amounting to 0.95 Md. A second plasmid (pAB524) with one *EcoRI* fragment (0.6 Md) of pAB124 deleted was also constructed. Restriction endonuclease cleavage maps of pAB224 and pAB524 were constructed.

### INTRODUCTION

Several plasmids have been isolated from Gram-positive spore-forming bacilli, e.g. *Bacillus pumilus* (Lovett & Bramucci, 1975), *B. megaterium* (Carlton & Helinski, 1969) and *B. subtilis* (Tanaka *et al.*, 1977; Le Hegarat & Anagnostopoulos, 1977), although none of these has been shown to be associated with any phenotypic trait. More recently, a plasmid from *B. cereus* coding for tetracycline resistance and one from *B. subtilis* producing a bacteriocin have been isolated (Bernhard *et al.*, 1978).

The presence of a plasmid in a bacteriocin (thermocin)-producing strain of *B. stearothermophilus* has recently been reported (Sharp *et al.*, 1979), and we have described the isolation of four plasmids from antibiotic-resistant thermophilic bacilli (Bingham *et al.*, 1979), one of which (pAB124) was shown to confer tetracycline resistance on *B. subtilis*. This paper describes the characterization of pAB124 and the construction of deletion variants of this plasmid.

### METHODS

**Bacterial strains.** Thermophilic *Bacillus* strain TB124, isolated as previously described (Bingham *et al.*, 1979), was classified as *Bacillus stearothermophilus* using standard identification procedures (Cowan & Steel, 1974). *Bacillus subtilis* harbouring plasmid pAB124 was prepared by direct plasmid transformation. *Bacillus subtilis* IG20 (*hsr hsm trp*) was kindly provided by A. Docherty, Department of Bacteriology, Bristol University.

**Culture media.** For plasmid isolation all strains were grown on TYS medium (Bingham *et al.*, 1979) supplemented with tetracycline at 25 µg ml<sup>-1</sup>. The following minimal medium (SMS) was used for *B. subtilis* (g l<sup>-1</sup>): (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 2; K<sub>2</sub>HPO<sub>4</sub>, 14; KH<sub>2</sub>PO<sub>4</sub>, 6; sodium citrate, 1; MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.2; glucose, 5.

**Isolation of plasmid DNA.** This was carried out with 400 ml cultures using the procedure previously described (Bingham *et al.*, 1979). Plasmid pAB124 was originally isolated from *B. stearothermophilus* TB124 and transformed into *B. subtilis* (Bingham *et al.*, 1979). All experiments described in this paper used pAB124 isolated from *B. subtilis*. Plasmid pUB1654 was a gift from A. Docherty.

† Present address: Department of Biochemistry, Imperial College, South Kensington, London SW7 2AZ.

**Small-scale plasmid preparation.** This was used for extracting recombinant DNA from large numbers of clones. Cultures (15 ml) in TYS medium containing the selective antibiotic(s) were grown overnight at 37 °C without shaking. Cells were harvested at 10000 g, washed once in 5 ml TES buffer (30 mM-Tris/HCl pH 8.0, 5 mM- $\text{Na}_2\text{EDTA}$ , 50 mM- $\text{NaCl}$ ) and resuspended in 100  $\mu\text{l}$  TES. EDTA (25  $\mu\text{l}$ , 0.25 M) and lysozyme (25  $\mu\text{l}$ , 15 mg  $\text{ml}^{-1}$ ) were added and the suspension was shaken gently on ice for 10 min. The suspension was then placed in a 37 °C water bath for 5 min and 25  $\mu\text{l}$  sodium dodecyl sulphate (10%, w/v) was added, to bring about lysis, followed by 50  $\mu\text{l}$   $\text{NaCl}$  (5 M); this mixture was left at 0 °C for 2.5 to 3 h. A cleared lysate was obtained by centrifugation at 45000 g for 45 min, and this was then extracted twice with chloroform/3-methylbutan-1-ol (24:1, v/v), once with phenol (freshly distilled over  $\text{Na}_2\text{O}$ ) equilibrated in TES buffer and three times with diethyl ether. The plasmid DNA obtained was suitable for restriction endonuclease analysis.

**Plasmid transformation.** *Bacillus subtilis* IG20 was grown overnight at 37 °C in 200 ml SMS medium supplemented with 0.5 ml tryptophan (2 mg  $\text{ml}^{-1}$ ) and then diluted with an equal volume of similar medium (prewarmed). The absorbance (540 nm) was monitored at 30 min intervals, and at a value equivalent to maximum competence (30 to 60 min after cessation of exponential growth) 1 ml samples of the culture were added to plasmid DNA (0.5 to 5  $\mu\text{g}$ ) in 50  $\mu\text{l}$  polyethylene glycol 1000 (50%, w/v) and incubated for 1 h at 37 °C with vigorous aeration. Prewarmed TYS medium (5 ml) was then added and incubation was continued for a further hour prior to plating on selective media. Selective concentrations of antibiotics used were 25  $\mu\text{g ml}^{-1}$  for tetracycline and neomycin and 75  $\mu\text{g ml}^{-1}$  for streptomycin. It was not necessary to induce competence by dilution of the culture into a starvation medium (Stacey, 1968).

**Restriction endonucleases.** *EcoRI* and *CauII* were purified by procedures developed in this laboratory and digestions were carried out in buffers previously described (Bingham *et al.*, 1979). *BstEII*, *HaeIII*, *HhaI*, *HindII*, *HpaI*, *HpaII*, *ThaI*, *XbaI* and T4-DNA ligase were purchased from Uniscience Ltd, Cambridge, and digestions carried out in the buffers recommended by the manufacturer. Reaction mixtures contained 0.5 to 1  $\mu\text{g}$  plasmid DNA in a final volume of 20 to 25  $\mu\text{l}$ , and reactions were terminated by adding  $\text{Na}_2\text{EDTA}$  to a final concentration of 10 mM (*CauII*, *BstEII*, *ThaI*) or heating at 66 °C for 10 min. For double digestions involving enzymes with different buffer requirements, plasmid DNA was first digested with the enzyme requiring the buffer of lower ionic strength, and then the buffer was adjusted with 10 $\times$  concentrated components prior to the addition of the second enzyme.

Agarose gels (0.8%, w/v) were used as previously described (Bingham *et al.*, 1979), with an *EcoRI* plus *HindIII*  $\lambda$  C1857 digest providing fragments of standard sizes: 13.4, 3.35, 3.2, 2.8, 2.32, 1.39, 1.27, 1.05, 0.89, 0.59, 0.47 and 0.31 megadaltons (Md) (Murray & Murray, 1975). The sizes of unknown fragments were determined graphically from a 10 $\times$  enlargement of a photograph (35 mm) and were taken as an average of three estimations.

**Cloning of pAB124 *EcoRI* fragments.** Plasmids pAB124 (1  $\mu\text{g}$ ) and pUB1654 (1  $\mu\text{g}$ ) were digested to completion with *EcoRI* and then the enzyme was denatured by heating at 66 °C for 10 min. The fragments were precipitated by adding 3 M-sodium acetate to a final concentration of 0.3 M and 2 vol. cold (-20 °C) ethanol. After 10 min in a solid  $\text{CO}_2$ /ethanol bath the precipitate was collected by centrifugation. After removal of all the ethanol the precipitate was resuspended in T4 ligase buffer (66 mM-Tris/HCl pH 7.6, 6.6 mM- $\text{MgCl}_2$ , 10 mM-dithiothreitol, 0.4 mM-ATP) and incubated overnight at 4 °C with 0.05 units T4-DNA ligase.

## RESULTS

### Restriction endonuclease site mapping of pAB124

The fragment sizes obtained by digestion of pAB124 DNA (isolated from *B. subtilis*) with *BstEII*, *CauII*, *EcoRI*, *HaeIII*, *HindII*, *HhaI*, *HpaI*, *HpaII*, *ThaI* and *XbaI* are given in Table 1. These sizes were determined graphically using a lambda DNA digested with *HindIII* plus *EcoRI* as standards subjected to electrophoresis on the same gels as the unknown fragments. The average value for the total size of pAB124 was close to 2.9 Md. A series of double digestion experiments was carried out in order to construct a cleavage map of pAB124 (Table 2).

The positions of the four single sites (*BstEII*, *CauII*, *HpaI*, *XbaI*) relative to each other were determined taking *CauII* as the reference point. An *HpaI* plus *CauII* digest and a *BstEII* plus *CauII* digest both gave a small fragment of 0.35 Md; thus the *BstEII* and *HpaI* sites are the same distance from *CauII* site either on the same side, i.e. very close together, or on opposite sides 0.7 Md apart. The latter position was confirmed by a *BstEII* plus *HpaI* double digestion. The *CauII* plus *XbaI* digestion generated fragments of 1.85 and 1.05 Md. It was assumed that the *XbaI* site was 1.85 Md to the right of *CauII* and the *HpaI*

*Bacillus-stearothermophilus* plasmids

111

Table 1. Fragments produced by restriction endonuclease digestion of pAB124

Digestions and agarose-gel electrophoresis were carried out as described in Methods using plasmid DNA isolated from *B. subtilis*.

Enzyme	Fragment size (Md)			Sum of fragment sizes (Md)
	A	B	C	
<i>Bst</i> EII	2.90			2.90
<i>Cau</i> II	2.90			2.90
<i>Hpa</i> I	2.90			2.90
<i>Xba</i> I	2.90			2.90
<i>Hha</i> I	2.23	0.70		2.93
<i>Hind</i> II	2.30	0.62		2.92
<i>Hae</i> III	1.50	1.18	0.25	2.93
<i>Hpa</i> II	1.80	0.70	0.42	2.92
<i>Tha</i> I	1.84	0.65	0.40	2.89
<i>Eco</i> RI	1.95	0.61	0.33	2.89

Table 2. Fragments produced by digestion of pAB124 with two restriction endonucleases

Double digestions and agarose-gel electrophoresis were carried out as described in Methods using plasmid DNA isolated from *B. subtilis*.

Enzyme pair	Fragment size (Md)				Sum of fragment sizes (Md)
	A	B	C	D	
<i>Bst</i> EII + <i>Cau</i> II	2.55	0.35			2.90
<i>Bst</i> EII + <i>Hpa</i> I	2.23	0.70			2.93
<i>Bst</i> EII + <i>Xba</i> I	2.20	0.70			2.90
<i>Cau</i> II + <i>Hpa</i> I	2.57	0.35			2.92
<i>Cau</i> II + <i>Xba</i> I	1.85	1.05			2.90
<i>Hpa</i> I + <i>Xba</i> I	1.48	1.40			2.88
<i>Hha</i> I + <i>Cau</i> II	2.20	0.44	0.25		2.89
<i>Hha</i> I + <i>Xba</i> I	1.59	0.71	0.60		2.90
<i>Hha</i> I + <i>Hpa</i> I	2.10	0.70	0.10		2.90
<i>Hind</i> II + <i>Cau</i> II	2.30	0.33	0.28		2.91
<i>Hind</i> II + <i>Xba</i> I	1.50	0.78	0.62		2.90
<i>Hind</i> II + <i>Hpa</i> I	2.30	0.61	—		2.91
<i>Hae</i> III + <i>Cau</i> II	1.24	1.18	0.25	0.24	2.91
<i>Hae</i> III + <i>Bst</i> EII	1.18	0.89	0.59	0.25	2.91
<i>Hae</i> III + <i>Xba</i> I	1.30	1.18	0.25	0.18	2.91
<i>Hae</i> III + <i>Hpa</i> I	1.51	1.19	0.16	0.10	2.96
<i>Hpa</i> II + <i>Cau</i> II	1.80	0.70	0.42	—	2.92
<i>Hpa</i> II + <i>Bst</i> EII	1.80	0.42	0.35	0.32	2.89
<i>Hpa</i> II + <i>Xba</i> I	1.80	0.70	0.38	(0.05)†	2.93
<i>Hpa</i> II + <i>Hpa</i> I	1.45	0.72	0.42	0.35	2.94
<i>Tha</i> I + <i>Cau</i> II	1.80	0.42	0.40	0.23	2.85
<i>Tha</i> I + <i>Bst</i> EII	1.83	0.62	0.40	(0.05)†	2.90
<i>Tha</i> I + <i>Xba</i> I	1.60	0.65	0.40	0.26	2.91
<i>Tha</i> I + <i>Hpa</i> I	1.75	0.65	0.40	0.10	2.90
<i>Eco</i> RI + <i>Cau</i> II	1.55	0.60	0.40	0.33	2.88
<i>Eco</i> RI + <i>Bst</i> EII	1.90	0.62	0.33	(0.05)†	2.90
<i>Eco</i> RI + <i>Xba</i> I	1.95	0.33*	0.29	—	2.88
<i>Eco</i> RI + <i>Hpa</i> I	1.20	0.75	0.60	0.32	2.87

\* This fragment was overabundant and assumed to be a 'double'.

† Fragment not detected, but its presence can be predicted.

and *Bst*EII sites were positioned using this orientation. If *Hpa*I lies 0.35 Md to the right of the *Cau*II site a double digestion with *Hpa*I plus *Xba*I would generate 1.48 and 1.40 Md fragments, whereas if it lies 0.35 Md to the left, 0.8 and 2.1 Md fragments would be generated. Since fragments of the former sizes were obtained, the *Hpa*I site must lie to the right of the




Fig. 1. Relative positions of fragments produced by digestion of pAB124 with several restriction endonucleases.

*CauII* site with the *BstEII* site 0.35 Md to the left. The positions of the four sites (Fig. 1) were consistent with the results of all the double digestions carried out.

Using these four single sites as reference points the fragments generated by *HhaI*, *HindII*, *HpaII*, *HaeIII*, *ThaI* and *EcoRI* (Table 1) were ordered using the established techniques of double-digestion mapping. For example, the *EcoRI* fragments were ordered easily as only one position of *EcoRI* fragment A was consistent with it containing the *BstEII*, *CauII* and *HpaI* sites but not the *XbaI* site which is present in *EcoRI* fragment B. A *CauII* plus *EcoRI* digestion generated 1.55 and 0.4 Md fragments from *EcoRI* A; the 1.55 Md fragment could only lie to the right of the *CauII* site with the 0.4 Md fragment to the left (Fig. 1). The *EcoRI* B fragment must lie to the right of *EcoRI* A over the *XbaI* site in the position shown (Fig. 1) since an *EcoRI* plus *XbaI* digest yielded 0.33 and 0.29 Md fragments from *EcoRI* B and the distance from *EcoRI* A to the *XbaI* site is 0.35 Md. The remaining fragment, *EcoRI* C, therefore lies to the right of *EcoRI* B.

The fragments generated by digestion of pAB124 with *HhaI*, *HindII*, *HaeIII*, *HpaII* and *ThaI* were ordered in a similar manner and their relative positions are shown in Fig. 1.

#### Cloning of *EcoRI* fragments of pAB124 carrying the tetracycline resistance gene

Staphylococcal plasmid pUB1654 conferring streptomycin and neomycin resistance was used. This has a single *EcoRI* site lying within the streptomycin resistance gene (A. Docherty, personal communication). It would therefore be expected that insertion of pAB124 fragments at this site would inactivate streptomycin resistance. Transformation of the ligated fragments generated about 35% of clones that were *Nm<sup>r</sup> Str<sup>r</sup> Tc<sup>r</sup>*. DNA was extracted from several of these clones and was analysed with *EcoRI* (Table 3). The minimum requirement for tetracycline resistance was shown to be the *EcoRI* fragment A (1.95 Md) of pAB124 present alone in 50% of the recombinant molecules examined.

#### Construction of small tetracycline resistance plasmids from cloned *EcoRI* fragments of pAB124

Plasmid DNA from each type of clone, representing all combinations of pAB124 *EcoRI* fragments (Table 3), was digested to completion with *EcoRI* and then ligated with T4-DNA ligase. The DNA was then transformed into *B. subtilis* IG20 and *Tc<sup>r</sup> Nm<sup>r</sup> Str<sup>r</sup>* clones were identified. Two types of plasmid were isolated from the clones examined (Table 4): pAB224

*Bacillus stearothermophilus* plasmids

113

Table 3. Restriction endonuclease analysis of recombinant plasmids of pAB124 and pUB1654

Plasmid DNA was extracted from clones of Nm<sup>r</sup> Str<sup>r</sup> Tc<sup>r</sup> phenotype and digested with *Eco*RI as described in Methods.

Clones identified		<i>Eco</i> RI digestion		Type of recombinant
No.	Plasmid size (Md)	Fragment size , (Md)	pAB124 fragment(s) inserted	
10	6.0	4.0*	—	I
		1.95	A	
3	6.55	4.0*	—	II
		1.95	A	
		0.60	B	
2	6.30	4.0*	—	III
		1.95	A	
		0.32	C	
4	6.90	4.0*	—	IV
		1.95	A	
		0.60	B	
		0.32	C	

\* This fragment corresponds to pUB1654.

Table 4. *Eco*RI analysis of tetracycline resistance plasmids derived from pAB124:pUB1654 recombinant molecules

Plasmid DNA was extracted from Tc<sup>r</sup> Nm<sup>r</sup> Str<sup>r</sup> clones and digested with *Eco*RI as described in Methods.

DNA source*		Tetracycline resistance plasmids			
Type of recombinant	pAB124 <i>Eco</i> RI fragment(s) present	No. of clones†	Size (Md)	<i>Eco</i> RI fragments (Md)	pAB124 <i>Eco</i> RI fragment(s) present
I	A	10	1.95	1.95	A (pAB224)
II	A + B	10	1.95	1.95	A (pAB224)
III	A + C	7	1.95	1.95	A (pAB224)
		3	2.3	1.95, 0.32	A + C (pAB524)
IV	A + B + C	8	1.95	1.95	A (pAB224)
		2	2.3	1.95, 0.32	A + C (pAB524)


\* Recombinant plasmids of pAB124 and pUB1654 (see Table 3).

† Ten clones from each transformation were examined.

which was the recircularized *Eco*RI fragment A of pAB124 (1.95 Md) and pAB524 containing *Eco*RI fragments A plus C (2.3 Md). No plasmids containing *Eco*RI fragments A plus B or reconstructed pAB124 were detected.

Plasmid pAB224 was shown to contain seven single sites (*Eco*RI, *Bst*EII, *Cau*II, *Hpa*II, *Hha*I, *Tha*I and *Hpa*I). The position of each site was determined by double digestions and was shown to be consistent with pAB224 being the recircularized *Eco*RI fragment A of pAB124 (Fig. 2). Therefore *Eco*RI A contains the tetracycline resistance genes and all the known essential functions of pAB124.


An *Eco*RI digest of pAB524 generated fragments of 1.95 Md and 0.32 Md. The larger fragment contained *Bst*EII, *Cau*II and *Hpa*I sites and double digestions confirmed that it was the *Eco*RI fragment A of pAB124; the smaller fragment contained *Hha*I and *Hpa*II sites and was assumed to be the *Eco*RI fragment C of pAB124. Two orientations of *Eco*RI C within pAB524 were possible and each would give different sized fragments when digested with *Hpa*II or *Hha*I. The observed sizes of the fragments obtained from five different examples of pAB524 indicated that the order of *Eco*RI fragments A and C was the same as that observed in pAB124 (Fig. 2).



## DISCUSSION

The development of the staphylococcal vector pUB1654 (Nm<sup>r</sup> Str<sup>r</sup>; A. Docherty, personal communication) allowed us to examine which region of pAB124 contained the

T. ATKINSON



124, pAB224 and pAB524.

pAB124 by standard procedures *stearothermophilus* but not identical is to develop a vector for cloning asis for such a vector. The expres- this plasmid as a vector for gene hat the plasmid is stable with no *btilis*, as determined by restriction. however, are lower with plasmid. rom *B. subtilis*; this may in part be nsform at a higher frequency than; venient restriction site for cloning; a cohesive terminus and lies within; tance or plasmid replication (Fig. ycyin resistance plasmid (pUB110); ce. Of the remaining single restric- *CauII* produce a cohesive terminus s making the use of these sites for; within the region of pAB124 that

654 (Nm<sup>+</sup> Str<sup>r</sup>; A. Docherty, per- region of pAB124 contained the

*Bacillus stearothermophilus* plasmids

115

tetracycline resistance genes. Only the large *EcoRI* fragment (A) was essential for the expression of the Tc<sup>r</sup> phenotype and the survival of the plasmid as a replicon. The circularized *EcoRI* fragment A (pAB224) contained one site for each of seven restriction endonucleases (Fig. 2); three of these enzymes produce a cohesive terminus (*EcoRI*, *HpaII* and *HhaI*), and therefore this plasmid is of potential use as a small (1.95 Md) vector for gene cloning. We have attempted to insert at the *EcoRI* site the *B. licheniformis*  $\beta$ -lactamase that is cloned between two *EcoRI* sites in a lambda vector (W.J. Brammar, personal communication) and also the  $\beta$ -lactamase of the staphylococcal plasmid pI258 (Novick *et al.*, 1979). In both cases we could not isolate recombinant molecules containing a  $\beta$ -lactamase gene. We are therefore continuing to examine the ability of pAB224 to express genes cloned at the *EcoRI* site.

The construction of pAB524 in a similar manner to pAB224 gives rise to a plasmid with two *EcoRI*, two *HpaII* and two *HhaI* sites, but no *XbaI* site. A potentially useful deletion plasmid would be that formed by circularization of *EcoRI* fragments A plus B of pAB124, which would retain the single *XbaI* site and also contain one *HhaI* site. However, repeated ligations of *EcoRI*-digested recombinants of pUB1654 containing *EcoRI* fragments A plus B failed to generate this plasmid; only pAB224 (*EcoRI* A) and pUB1654 containing *EcoRI* fragment A of pAB124 were obtained. This was rather surprising since one would expect circularization of *EcoRI* fragments A plus B to be about as efficient as circularization of *EcoRI* fragments A plus C (pAB524; Table 4).

We intend to examine the ability of pAB124 and pAB224 to express chromosomal functions cloned at the *XbaI* and *EcoRI* sites, respectively, in order to evaluate their use as vectors for gene cloning in *Bacillus* species.

We thank A. Docherty for providing pUB1654 DNA. This work was supported by a grant from the Health and Safety Executive.

## REFERENCES

- BERNHARD, K., SCHREMPF, H. & GOEBEL, W. (1978). Bacteriocin and antibiotic resistant plasmids in *Bacillus cereus* and *Bacillus subtilis*. *Journal of Bacteriology* 133, 897-903.
- BINGHAM, A. H. A., BRUTON, C. J. & ATKINSON, T. (1979). Isolation and partial characterization of four plasmids from antibiotic-resistant thermophilic bacilli. *Journal of General Microbiology* 114, 401-408.
- CANOSI, U., MORELLI, G. & TRAUTNER, T. A. (1978). The relationship between molecular structure and transformation efficiency of some *S. aureus* plasmids isolated from *B. subtilis*. *Molecular and General Genetics* 166, 259-267.
- CARLTON, B. C. & HELINSKI, D. R. (1969). Heterogeneous circular DNA elements in vegetative cultures of *Bacillus megaterium*. *Proceedings of the National Academy of Sciences of the United States of America* 65, 592-599.
- COWAN, S. T. & STEEL, K. J. (1974). *Identification of Medical Bacteria*, 2nd edn. Cambridge: Cambridge University Press.
- LE HEGARAT, J.-C. & ANAGNOSTOPOULOS, C. (1977). Detection and characterisation of naturally occurring plasmids in *Bacillus subtilis*. *Molecular and General Genetics* 157, 167-174.
- LOVETT, P. S. & BRAMUCCI, M. G. (1975). Plasmid deoxyribonucleic acid in *Bacillus pumilus* and *Bacillus subtilis*. *Journal of Bacteriology* 124, 484-490.
- MURRAY, K. & MURRAY, N. E. (1975). Phage lambda receptor chromosome for DNA fragments with *HindIII* and *EcoRI*. *Journal of Molecular Biology* 98, 551-564.
- NOVICK, R. P., MURPHY, E., GRYCZAN, T. J., BARON, E. & EDELMAN, I. (1979). Penicillinase plasmids of *Staphylococcus aureus*: restriction-deletion maps. *Plasmid* 2, 109-129.
- SHARP, R. J., BINGHAM, A. H. A., COMER, M. J. & ATKINSON, A. (1979). Partial characterization of a bacteriocin (thermocin) from *Bacillus stearothermophilus* RS93. *Journal of General Microbiology* 111, 449-451.
- STACEY, K. A. (1968). Synthesis and properties of DNA: transformation. In *Experiments in Microbial Genetics*, pp. 46-51. Edited by R. C. Clowes & W. Hayes. Oxford: Blackwell Scientific Publications.
- TANAKA, T., KURODA, M. & SAKAGUCHI, K. (1977). Isolation and characterisation of four plasmids from *Bacillus subtilis*. *Journal of Bacteriology* 129, 1487-1494.





CGCTCAGGATTCGACTTCGCGCTAGAGGATCGGATCCCGGCGCGATTATATAGCTCGATC  
 TTCTCTATATTCGCGCTAGAGGATCGGATCCCGGCGCGATTATATAGCTCGATC  
 CGCTCAGGATTCGACTTCGCGCTAGAGGATCGGATCCCGGCGCGATTATATAGCTCGATC  
 TTCTCTATATTCGCGCTAGAGGATCGGATCCCGGCGCGATTATATAGCTCGATC

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

Search

Nucleotide

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

Display

default


Show:

20

Send to

File

Get Subsequence


☐ 1: L09136. pUC18c cloning ve...[gi:209210]

Links

LOCUS SYNPU18CV 2686 bp DNA circular SYN 27-APR-1993  
 DEFINITION pUC18c cloning vector (beta-galactosidase mRNA on complementary strand).  
 ACCESSION L09136  
 VERSION L09136.1 GI:209210  
 KEYWORDS beta-galactosidase; cloning vector.  
 SOURCE synthetic construct  
 ORGANISM synthetic construct  
 artificial sequences.  
 REFERENCE 1 (bases 1 to 2686)  
 AUTHORS Gilbert,W.  
 TITLE Obtained from VecBase 3.0  
 JOURNAL Unpublished (1991)  
 COMMENT Original source text: Synthetic construct DNA.  
 These data and their annotation were supplied to GenBank by Will Gilbert under the auspices of the GenBank Curator Program. pUC18c - Cloning vector (beta-galactosidase mRNA on complementary strand)  
 ENTRY PUC18C #TYPE DNA CIRCULAR TITLE pUC18c - Cloning vector (beta-galactosidase mRNA on complementary strand)  
 DATE 02-APR-1986  
 #sequence 16-DEC-1986  
 ACCESSION VB0032  
 SOURCE artificial  
 COLLECTION ATCC 37253  
 REFERENCE  
 #number 1  
 #authors Norrander J., Kempe T., Messing J.  
 #journal Gene (1983) 26: 101-106  
 REFERENCE  
 #number 2  
 #authors Pouwels P.H., Enger-Valk B.E., Brammar W.J.  
 #book Cloning Vectors, Elsevier 1985 and supplements  
 #comment vector I-A-iv-20  
 COMMENT  
 Assembled from pUC19c and M13mp18 by F. Pfeiffer, MPI, Martinsried  
 Revised 16-DEC-1986 by F. Pfeiffer:  
 1062/3 'AT' to 'TA' to match revised sequence of PBR322  
 The beta-galactosidase mRNA sequence including the multiple cloning site of M13mp18 is on the strand complementary to that shown.  
 KEYWORDS  
 CROSSREFERENCE  
 #complement  
 VecBase(3):pUC18  
 PARENT



## Features of pUC18c (2686 bp)

residue	source
1- 137	2074-2210 pBR322
138- 237	2252-2351 pBR322
238- 396	1461-1303 (c) Lac-Operon
399- 455	57- 1 (c) polylinker of M13mp18
456- 682	1297-1071 (c) Lac-Operon
683-2686	2352-4355 pBR322

## Conflict (cfl) and Mutations (mut):

pUC18c	source
mut 1942 A G 3611	pBR322
mut 2243 T C 3912	pBR322

## FEATURE

1629-2417 789-1 (c) Ap-R; b-lactamase

POLYLINKER EcoRI-SacI-KpnI-SmaI-BamHI-XbaI-SalI-PstI-SphI-HindIII

## SELECTION

#resistance Ap

#indicator beta-galactosidase

SUMMARY pUC18c #length 2686 #checksum 3662.

## FEATURES

source

Location/Qualifiers

1..2686

/organism="synthetic construct"

/db\_xref="taxon:32630"

## BASE COUNT

666 a 677 c 684 g 659 t

## ORIGIN

```

1  tcgcgcgttt  cggatgatgac  ggtgaaaacc  tctgacacat  gcagctcccg  gagacgggtca
61  cagcttgtct  gtaagcggat  gccgggagca  gacaagcccc  tcagggcgcg  tcagcgggtg
121  ttggcgggtg  tcggggctgg  cttaactatg  cggtatcaga  gcagattgta  ctgagagtgc
181  accatatgcg  gtgtgaaata  ccgcacagat  gcgtaaggag  aaaataccgc  atcaggcgcc
241  attcgccatt  caggctgctg  aactgttggg  aagggcgatc  ggtgcgggcc  tcttcgctat
301  tacgccagct  ggcgaaagg  ggatgtgctg  caaggcgatt  aagttgggta  acgccagggt
361  tttcccagtc  acgacgttgt  aaaacgacgg  ccagtgccaa  gcttgcatgc  ctgcaggctcg
421  actctagagg  atccccgggt  accgagctcg  aattcgtaat  catggtcata  gctgtttcct
481  gtgtgaaatt  gttatccgct  cacaattcca  cacaacatac  gagccggaag  cataaagtgt
541  aaagcctggg  gtgcctaatt  agtgagctaa  ctcacattaa  ttgcgttgcg  ctcactgccc
601  gctttccagt  cgggaaacct  gtcgtgccag  ctgcattaa  gaatcgcca  acgcgcgggg
661  agaggcgggt  tgcgtattgg  gcgctcttcc  gcttcctcgc  tctactgactc  gctgcgctcg
721  gtcgttcggc  tgcggcgagc  ggtatcagct  cactcaaagg  cggtaatagc  gttatccaca
781  gaatcagggg  ataacgcagg  aaagaacatg  tgagcaaaa  gccagcaaaa  ggccaggaa
841  cgtaaaaagg  ccgctgtgct  ggcgttttcc  cataggctcc  gccccctga  cgagcatcac
901  aaaaaatgac  gctcaagtca  gaggtggcga  aacccgacag  gactataaag  ataccaggcg
961  tttccccctg  gaagctccct  cgtgcgctct  cctgttccga  ccctgccgct  taccggatac
1021  ctgtccgctt  ttctcccttc  gggaagcgtg  gcgctttctc  atagctcacg  ctgtagggtat
1081  ctgagttcgg  tgtaggctcg  tcgctccaag  ctgggctgtg  tgcacgaacc  ccccggttcag
1141  cccgaccgct  gcgccttacc  cggttaactat  cgtcttgagt  ccaacccggt  aagacacgac
1201  ttatcgccac  tggcagcagc  cactggtaac  aggattagca  gagcgaggta  tgtaggcggg
1261  gctacagagt  tcttgaagtg  gtggcctaac  tacggctaca  ctagaaggac  agtatttggt
1321  atctgcgctc  tgctgaagcc  agttaccttc  ggaaaaagag  ttggtagctc  ttgatccggc
1381  aaacaaacca  ccgctggtag  cgggtggttt  tttgtttgca  agcagcagat  tatgcgcaga
1441  aaaaaaggat  ctcaagaaga  tcctttgatc  ttttctacgg  ggtctgacgc  tcagtggaac
1501  gaaaactcac  gttaagggat  tttgggtcatg  agattatcaa  aaaggatctt  caccatagatc
1561  ctttttaaatt  aaaaatgaag  ttttaaatca  atctaaaagta  tatatgagta  aacttggtct
1621  gacagttacc  aatgcttaat  cagtgaggca  cctatctcag  cgatctgtct  atttcgttca
1681  tccatagttg  cctgactccc  cgtcgtgtag  ataactacga  tacgggaggg  cttaccatct
1741  ggccccagtc  ctgcaatgat  accgcgagac  ccacgctcac  cggctccaga  tttatcagca
1801  ataaaccagc  cagccggaag  ggccgagcgc  agaagtgggtc  ctgcaacttt  atccgcctcc
1861  atccagtcta  ttaattgttg  ccgggaagct  agagtaagta  gttcgccagt  taatagtttg
1921  cgcaacgttg  ttgccattgc  tacaggcatc  gtggtgtcac  gctcgtcgtt  tgggtatggc
1981  tcattcagct  ccggttccca  acgatcaagg  cgagttacat  gatcccccat  gttgtgcaaa
2041  aaagcgggta  gctccttcgg  tcctccgac  gttgtcagaa  gtaagtggc  cgcagtggtta

```


```
2101 tcactcatgg ttatggcagc actgcataat tctcttactg tcatgccatc cgtaagatgc
2161 ttttctgtga ctggtgagta ctcaaccaag tcattctgag aatagtgtat gcggcgaccg
2221 agttgctctt gcccggcgtc aatacgggat aataccgcgc cacatagcag aactttaaaa
2281 gtgctcatca ttggaaaacg ttcttcgggg cgaaaactct caaggatctt accgctgttg
2341 agatccagtt cgatgtaacc cactcgtgca cccaactgat cttcagcatc ttttactttc
2401 accagcgttt ctgggtgagc aaaaacagga aggcaaaatg ccgcaaaaaa gggaataagg
2461 gcgacacgga aatggttgaat actcatactc ttcctttttc aatattattg aagcatttat
2521 caggggttatt gtctcatgag cggatacata tttgaatgta tttagaaaaa taaacaaata
2581 ggggttcgc gcacatttcc ccgaaaagtg ccacctgacg tctaagaaac cattattatc
2641 atgacattaa cctataaaaa taggcgtatc acgaggccct ttcgtc
```

//

Revised: July 5, 2002.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

Jan 21 2003 18:08:12



## PATENT

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

<i>In re</i> Application of	)	
E. GREEN <i>et al.</i>	)	Group Art Unit: 1652
Serial No. 09/754,083	)	Examiner: M. N. RAO
Filed: January 5, 2001	)	Atty. Docket No. 000487.00007
For: <b>ETHEANOL PRODUCTION</b>	)	

DECLARATION

Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

We, Edward Green, Muhammad Javed, and Namdar Baghaei-Yazdi, declare that:

1. We are the named inventors of the subject matter claimed in the above-identified application. We deposited *Bacillus* strains LN (accession number 41038), TN (accession number 41039), LN-S (J8) (accession number 41040), and LN-D (accession number 41041) with the National Collections of Industrial, Food and Marine Bacteria (NCIMB) at 23 St. Machar Drive, Aberdeen AB2 1RY, Scotland, United Kingdom. Each deposit was made under the terms of the Budapest treaty.

2. Upon issuance of a patent each of these strains will be irrevocably and without restriction or conditions be available to the public.

3. We hereby declare that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these were made with the knowledge that false statements made willfully are punishable by fine, imprisonment, or both a fine and imprisonment under Section 1001 of Title 18 of the United States; and further that false statements made willfully may jeopardize the validity of any patent issuing on an application in which the false statements were made.

20/01/03

Date

17/01/03

Date

17/1/03

Date

EMC

Edward Green

Muhammad Javed

Muhammad Javed

Namdar Baghaei

Namdar Baghaei-Yazdi

FEMS Microbiology Letters 26 (1985) 333-336  
Published by Elsevier

FEM 02008

333

EXHIBIT

## Mutants of *Bacillus stearothermophilus* lacking NAD-linked L-lactate dehydrogenase

(Fermentation; ethanol production; fluoropyruvate resistance; thermophile)

Mark A. Payton and Brian S. Hartley \*

Department of Microbiology, Biogen S.A., P.O. Box 1211, Geneva 24, Switzerland, and \* Centre for Biotechnology, Imperial College,  
London SW7 2AZ, U.K.

Received 15 November 1984

Accepted 23 November 1984

### 1. SUMMARY

*Bacillus stearothermophilus* produces low levels of ethanol following fermentation of glucose at temperatures in excess of 60°C. A major by-product is L-lactic acid. To improve ethanol yields mutants of this organism were isolated which lacked NAD-linked L-lactate dehydrogenase, the enzyme responsible for lactic acid production. This was achieved by selection for resistance to a suicide substrate, fluoropyruvate. Such mutants no longer produced lactic acid and yields of ethanol from glucose were increased by at least a factor of two.

### 2. INTRODUCTION

*B. stearothermophilus* is a facultatively anaerobic thermophilic bacillus capable of growing either oxidatively or fermentatively on glucose [1]. During fermentative growth in rich broth with glucose or sucrose as the major carbon source *B. stearothermophilus* displays a "mixed-acid" type fermentation with the production of lactic, acetic and formic acids and some ethanol [2].

The oxidation and fermentative metabolism of glucose by wild-type *B. stearothermophilus*

(NCA1503) has been described previously [3,4]. It appears similar to that typified by the enteric bacterium *Escherichia coli* in that under aerobic, oxidative conditions, pyruvate formed by glycolysis is further metabolised by a pyruvate dehydrogenase (PDH) complex [5] to form acetyl CoA, which would normally be oxidised through the citric acid cycle. Growth under fermentative conditions, however, induces the synthesis of a pyruvate formate lyase (PFL) system [6] which converts the pyruvate to acetyl CoA, but with the concomitant production not of NADH as with the PDH complex but of formate. It has been reported previously both for *B. stearothermophilus* itself [2] and for other "mixed-acid" type fermentative microorganisms such as *E. coli* [7] that the pH of a fermentation can drastically affect the ratios of the different possible end-products. For *B. stearothermophilus*, there appears to be a relationship whereby the production of low levels of lactic acid as a fermentative end-product facilitates the production of relatively higher amounts of ethanol. This is due, presumably, to the extra availability both of carbon as pyruvate and of reducing equivalents as NADH under such conditions. We have been investigating the effects of a variety of parameters on the overall fermentation balance of

this organism as a model system for the improvement of ethanol-producing thermophiles for possible use in a thermophilic ethanol fermentation process. In order to improve ethanol yields, therefore, we attempted to abolish completely the production of lactic acid during the fermentation of glucose by *B. stearothermophilus* by the manipulation of media and growth conditions, but without total success. We therefore opted to remove NAD-linked lactate dehydrogenase, the enzyme responsible for L-lactate production, by mutagenesis. We describe here the isolation of mutants lacking this enzyme and a preliminary discussion of their fermentation characteristics.

### 3. MATERIALS AND METHODS

The starting strain of *B. stearothermophilus* used in this study was a prototrophic variant of *B. stearothermophilus* NCA1503. This strain, PSII, was isolated previously as a strain capable of rapid anaerobic growth on glucose (Payton, unpublished). Cultures were grown and maintained on LB glucose medium containing yeast extract (20 g · l<sup>-1</sup>), tryptone (10 g · l<sup>-1</sup>) and D-glucose (2 g · l<sup>-1</sup>) at a final pH of 7.0. For plates, LB glucose medium was solidified with 1.5% (w/v) Difco Bacto Agar. Growth was performed at 60°C unless otherwise stated.

Fluoropyruvic acid (sodium salt) was obtained from Sigma (London). Mutants of *B. stearothermophilus* PSII resistant to this compound were isolated first by growing PSII overnight at 60°C aerobically in LB glucose medium. Samples of 0.1 ml of these cultures containing approximately 10<sup>8</sup> viable cells were then plated onto LB glucose medium solidified with agar and allowed to dry. Fluoropyruvate was then applied, as a solid (2–5 mg) to the centre of each plate and the plates were incubated overnight at 60°C, after which time zones of inhibition of 1–3 cm could be seen within a lawn of bacterial growth. Occasionally single colonies could be seen within the zone of inhibition and these were picked onto LB glucose solid medium for purification. Single colonies purified in this way were retested by repeating the original

isolation procedure. Only those strains which exhibited zones of inhibition repeatedly and significantly less than the parental strain PSII (i.e. < 0.5 cm) were chosen for further study.

Initial classification of these strains was performed by growing the strains individually in 2-ml volumes of LB glucose medium in static culture at 60°C for 12 h. Half of the culture was centrifuged at 5000 × g for 5 min and the resultant supernatant assayed for the presence of L-lactic acid using a commercially available kit for the determination of L-lactic acid (Boehringer). The remaining half of the culture was permeabilised by a modification of the method previously published for *E. coli* [8].

To 1 ml samples of culture, 10 µl aliquots of a 10% (v/v) solution of toluene in absolute ethanol were added. Following incubation on ice for 20 min the mixtures were vortexed for 1 min and returned to ice. Samples were used within 30 min of preparation to measure NAD-linked L-lactate dehydrogenase by the method of Tarmy and Kaplan [9].

The assay contained in a total volume of 1 ml: NADH (0.3 mM), sodium pyruvate (30 mM) in potassium phosphate buffer (0.1 M, pH 7.5). Sufficient permeabilised extract was added to give an absorbance change of approx. 0.1 min<sup>-1</sup> at 340 nm and 25°C. Assay results were verified by preparing sonicates of the appropriate strain; cells from 100 ml overnight cultures in LB glucose medium were harvested by centrifugation, washed twice in 10 ml volumes of potassium phosphate buffer (0.1 M, pH 7.5) and finally resuspended in 5 ml of the same buffer. Suspensions were sonicated for 5 1-min bursts (150 W MSE ultrasonic disintegrator, peak to peak amplitude 12 µm) at 0°C and centrifuged at 4°C and 12000 × g for 15 min to remove debris. Lactate dehydrogenase was then assayed in supernatants as described above.

Determinations of the products of glucose fermentation were performed on supernatants from cultures grown on either CG medium (tryptone 30 g · l<sup>-1</sup>, yeast extract 10 g · l<sup>-1</sup>, D-glucose 10 g · l<sup>-1</sup>) or SG medium (tryptone 1 g · l<sup>-1</sup>, yeast extract 0.5 g · l<sup>-1</sup>, D-glucose 10 g · l<sup>-1</sup>) for 40 h at 57°C. Assays for lactic acid were performed as described

as which ex-  
and signifi-  
III (i.e. < 0.5

ns was per-  
ally in 2-ml  
tic culture at  
centrifuged  
ultant super-  
L-lactic acid  
for the de-  
ger). The re-  
abilised by a  
ly published

aliquots of a  
olute ethanol  
n ice for 20  
1 min and  
within 30 min  
ked L-lactate  
Tarmy and

ume of 1 ml:  
(30 mM) in  
H 7.5). Suffi-  
to give an  
nin<sup>-1</sup> at 340  
ified by pre-  
strain; cells

LB glucose  
ation, washed  
m phosphate  
suspended in  
as were soni-  
SE ultrasonic  
le 12 µm) at  
00 × g for 15  
rogenase was  
ribed above.  
f glucose fer-  
natants from  
(tryptone 30  
ose 10 g · l<sup>-1</sup>)  
yeast extract  
30 h at 57°C.  
as described

above. Ethanol and acetate were also measured using commercially available kits from Sigma (U.K.) and Boehringer, respectively.

#### 4. RESULTS AND DISCUSSION

We had previously failed to isolate lactate dehydrogenase-deficient strains by conventional techniques such as screening for low acid-producing mutants of *B. stearothersophilus* PSII following growth on glucose either aerobically or anaerobically. We therefore sought an alternative method based on a selection for resistance to a suicide substrate, fluoropyruvate. We had previously noted that the pyruvate analogue fluoropyruvate (FCH<sub>2</sub>-CO-COOH) was toxic to *B. stearothersophilus*. The toxicity of fluoropyruvate could depend on its ability itself to inhibit a key metabolic step, or could depend on its conversion initially either to fluoroacetyl CoA or to fluorolactate. In each case, amongst fluoropyruvate-resistant strains, some should be defective in uptake or accumulation of the toxic analogue. In addition, however, mutants should be isolated which affect the target reaction of inhibition or the metabolism of fluoropyruvate to fluoroacetyl CoA (via PDH or PFL) or to fluorolactate (via NAD-linked L-lactate dehydrogenase).

We therefore isolated fluoropyruvate-resistant mutants (MATERIALS AND METHODS) and proceeded to characterise the resulting strains. Strains chosen for further study, designated FPY (fluoropyruvate-resistant) mutants, were analysed by screening glucose-grown cells for the presence or absence of NAD-linked L-lactate dehydrogenase. This was tested initially on cells permeabilised with toluene, then verified using cell extracts prepared by sonication (MATERIALS AND METHODS). The supernatants from the same cultures were also assayed for the presence of L-lactic acid (MATERIALS AND METHODS). The results of these analyses are shown in Table 1. Of 12 strains thus tested 8 failed to produce lactic acid and also lacked detectable LDH activity in permeabilised cell suspensions. Assays performed on sonicates of selected strains were used to verify the absence of LDH. One of these strains, FPY 15, was chosen for

Table 1

An analysis of fluoropyruvate-resistant mutants of *B. stearothersophilus* PSII

Strain No.	Lactate production	LDH detected in toluenised cells	LDH present in sonicates
FPY 4	+	+	+
FPY 5	-	-	-
FPY 6	-	-	-
FPY 7	+	+	+
FPY 8	+	+	NT*
FPY 9	+	+	NT
FPY 10	-	Low	NT
FPY 11	-	Low	NT
FPY 12	-	-	NT
FPY 13	-	Low	NT
FPY 14	-	-	NT
FPY 15	-	-	-

\* NT, not tested.

further study and was redesignated *Ild15* (L-lactate dehydrogenase-deficient) and its fermentation balance studied.

Table 2 shows a comparison of the major fermentation products from glucose of *Ild15* vs. its parent PSII. First, in both strains acetate formation is higher on complex medium than on simple medium. We believe this is due to conversion of amino acids in the complex medium into acetate. Second, ethanol production by *Ild15* is higher by a factor of two than by PSII. We have subsequently performed a number of fermentations on *Ild15* both in batch and continuous culture at varying pH which have been published elsewhere [3,4] and have improved the ethanol yield by *Ild15*. The figure quoted for PSII of 0.5 mol ethanol per

Table 2

Yields of fermentation products from glucose (mol product/mol glucose) for *B. stearothersophilus* strains PSII and *Ild15*

The media used and assays of products are described in MATERIALS AND METHODS.

Strain	PSII		<i>Ild15</i>	
	CG	SG	CG	SG
Medium				
Lactate	1	1	0.04	0
Acetate	2	0.5	3.38	1
Ethanol	0.5	0.5	1	1
Total	3.5	2.0	4.42	2.0

mol glucose is, however, the highest we have obtained to date; at lower pHs ethanol production drops off sharply. These data show first that mutants of *B. stearothersophilus* lacking NAD-linked lactate dehydrogenase retain the ability to grow on glucose aerobically or anaerobically. Furthermore, the lactate dehydrogenase-deficient (*Ild*) strains of *B. stearothersophilus* produce significantly more ethanol than the wild-type strain following fermentation of glucose, verifying the trends shown by McKray and Vaughn [2]. Results from fermentations performed on *Ild15* suggest that the isolation of similar mutants in other organisms of potential use in ethanol fermentations may be a means of increasing ethanol yields. In addition, the isolation of *Ild* mutants of *B. stearothersophilus* amongst fluoropyruvate-resistant strains suggests that in this organism one mode of fluoropyruvate toxicity is dependent on its initial conversion to fluorolactate, in contrast to previous reports of the effect of this and similar fluoroanalogues on *E. coli* [10]. We have, in fact, shown (Payton, M.A., unpublished) that fluoropyruvate is a substrate, albeit a poor one, for semipurified LDH from *B. stearothersophilus*, as has been reported for LDHs from other sources [11].

## REFERENCES

- [1] Atkinson, A., Ellwood, D.C., Evans, C.G.T. and Yeo, R.G. (1975) *Biotech. Bioeng.* 17, 1375-1377.
- [2] McKray, G.A. and Vaughn, R.H. (1957) *Food Res.* 22, 494-500.
- [3] Hartley, B.S. and Payton, M.A. (1983) *Biochem. Soc. Symp.* 48, 133-146.
- [4] Hartley, B.S., Payton, M.A., Pyle, D.L., Mistry, P. and Shama, G. (1983) in *Biotech* 83, pp. 895-905. Online, London, U.K.
- [5] Henning, U., Dennert, G., Hertel, R. and Shipp, W.S. (1966) *Cold Spring Harbor Symp. Quant. Biol.* 31, 227-234.
- [6] Knappe, J., Blaschkowski, H.P., Grobner, P. and Schmitt, T. (1974) *Eur. J. Biochem.* 50, 253-263.
- [7] Dawes, E.A. and Foster, S.M. (1956) *Biochim. Biophys. Acta* 22, 253-265.
- [8] Reeves, R.E. and Sols, A. (1973) *Biochem. Biophys. Res. Commun.* 50, 459-466.
- [9] Tarmy, E.M. and Kaplan, N.O. (1968) *J. Biol. Chem.* 243, 2579-2586.
- [10] Avi-Dor, Y. and Mager, J. (1956) *J. Biol. Chem.* 222, 249-258.
- [11] Eisman, E.H., Lee Jr., J.A. and Winer, A.D. (1965) *Biochemistry* 4, 606-610.